antibodies -online.com





COMP Protein (Myc-DYKDDDDK Tag)



Image



$C \circ + \circ$	Drad	110+	2000
Go to	riou	uCt	Daue

()	11	\sim	rv		۱ ۸
	1 \ /	┙	I \/	╙	1/1

KDDDDK Tag.
essed in HEK293 cells.
omassie blue staining
ollagenous extracellular matrix (ECM) protein. It nits, each with EGF-like and calcium-binding ration results from formation of a five-stranded ECM proteins such as collagen appears to depend on of a 5 aa aspartate repeat and other mutations can

Target Details

	cause pseudochondroplasia (PSACH) and multiple epiphyseal dysplasia (MED).	
Molecular Weight:	80.9 kDa	
NCBI Accession:	NP_000086	


Application Details

Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only

Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.

Images



Western Blotting

Image 1. Validation with Western Blot