# antibodies - online.com







## CSNK1G3 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Overview

#### Image



| 0 - 1 - | DI.    | 4     |      |
|---------|--------|-------|------|
| L-O TO  | ₽r∩ai  | ICT I | nane |
| Go to   | 1 1001 | uct   | page |

| Quantity:                     | 20 μg   |
|-------------------------------|---|
| Target:                       | CSNK1G3   |
| Protein Characteristics:      | Transcript Variant 1                                    |
| Origin:                       | Human   |
| Source:                       | HEK-293 Cells   |
| Protein Type:                 | Recombinant   |
| Purification tag / Conjugate: | This CSNK1G3 protein is labelled with Myc-DYKDDDDK Tag. |
| Application:                  | Antibody Production (AbP), Standard (STD)               |

#### **Product Details**

| Characteristics: | <ul> <li>Recombinant human CSNK1G3 (transcript variant 1) protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul> |
|------------------|--|
| Purity:          | > 80 % as determined by SDS-PAGE and Coomassie blue staining   |
| Target Details   |  |

| Target:           | CSNK1G3  |
|-------------------|--|
| Alternative Name: | Csnk1g3 (CSNK1G3 Products)   |
| Background:       | This gene encodes a member of a family of serine/threonine protein kinases that                  |
|                   | phosphorylate caseins and other acidic proteins. A related protein in the African clawed frog    |
|                   | participates in the transmission of Wnt/beta-catenin signaling. Alternatively spliced transcript |
|                   | variants encoding multiple isoforms have been observed for this gene.                            |

#### Target Details

| Molecular Weight: | 51.2 kDa           |
|-------------------|--------------------|
| NCBI Accession:   | NP_004375          |
| Pathways:         | Hedgehog Signaling |


### **Application Details**

| Application Notes: | Recombinant human proteins can be used for:          |
|--------------------|--|
|                    | Native antigens for optimized antibody production    |
|                    | Positive controls in ELISA and other antibody assays |
| Comment:           | The tag is located at the C-terminal.                |
| Restrictions:      | For Research Use only                                |

#### Handling

| Concentration:   | 50 μg/mL  |
|------------------|---|
| Buffer:          | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.  |
| Storage:         | -80 °C  |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

#### Images



#### **Western Blotting**

Image 1. Validation with Western Blot