antibodies -online.com







CYB5R3 Protein (Transcript Variant 2) (Myc-DYKDDDDK Tag)



Image




| 0 - 1 - | DI. | 4 | |
|---------|-------|-------|------|
| Go to | Prodi | ICT r | nane |
| 00 10 | Hout | iot p | Jayc |

| () | ve | K\ / | | A . |
|-----|--------|------|----|-----|
| | \cup | 1 V/ | Щ. | V۷ |
| | | | | |

| Overview | |
|-------------------------------|---|
| Quantity: | 20 μg |
| Target: | CYB5R3 |
| Protein Characteristics: | Transcript Variant 2 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This CYB5R3 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |
| Product Details | |
| Characteristics: | Recombinant human CYB5R3 (transcript variant 2) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| Target Details | |
| Target: | CYB5R3 |
| Alternative Name: | Cyb5r3 (CYB5R3 Products) |
| Background: | This gene encodes cytochrome b5 reductase, which includes a membrane-bound form in |
| | somatic cells (anchored in the endoplasmic reticulum, mitochondrial and other membranes) |
| | and a soluble form in erythrocytes. The membrane-bound form exists mainly on the |
| | cytoplasmic side of the endoplasmic reticulum and functions in desaturation and elongation of |

| Target Details | |
|---------------------|---|
| | fatty acids, in cholesterol biosynthesis, and in drug metabolism. The erythrocyte form is located in a soluble fraction of circulating erythrocytes and is involved in methemoglobin reduction. The membrane-bound form has both membrane-binding and catalytic domains, while the soluble form has only the catalytic domain. Alternate splicing results in multiple transcript variants. Mutations in this gene cause methemoglobinemias. |
| Molecular Weight: | 31.4 kDa |
| NCBI Accession: | NP_015565 |
| Pathways: | SARS-CoV-2 Protein Interactome |
| Application Details | |
| Application Notes: | Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |
| Handling | |
| Concentration: | 50 μg/mL |

| Concentration: | 50 μg/mL | |
|------------------|--|--|
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. | |
| Storage: | -80 °C | |
| Storage Comment: | orage Comment: Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. | |



Western Blotting

Image 1. Validation with Western Blot