antibodies -online.com





COASY Protein (Transcript Variant 5) (Myc-DYKDDDDK Tag)



Image



Go to Product page

Overview

OVEIVIEVV	
Quantity:	20 μg
Target:	COASY
Protein Characteristics:	Transcript Variant 5
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This COASY protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	 Recombinant human Coenzyme A synthase (COASY), nuclear gene encoding mitochondrial protein, transcript variant 5 (transcript variant 5) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	COASY
Alternative Name:	Coenzyme A Synthase (Coasy) (COASY Products)
Background:	Coenzyme A (CoA) functions as a carrier of acetyl and acyl groups in cells and thus plays an important role in numerous synthetic and degradative metabolic pathways in all organisms. In

eukaryotes, CoA and its derivatives are also involved in membrane trafficking and signal

transduction. This gene encodes the bifunctional protein coenzyme A synthase (CoAsy) which carries out the last two steps in the biosynthesis of CoA from pantothenic acid (vitamin B5). The phosphopantetheine adenylyltransferase domain of this bifunctional protein catalyzes the conversion of 4'-phosphopantetheine into dephospho-coenzyme A (dpCoA) while its dephospho-CoA kinase domain completes the final step by phosphorylating dpCoA to form CoA. Mutations in this gene are associated with neurodegeneration with brain iron accumulation (NBIA). Alternative splicing results in multiple isoforms.


Molecular Weight:	30 kDa
NCBI Accession:	NP_001035996
Pathways:	Ribonucleoside Biosynthetic Process

Application Details

Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only

Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot