antibodies -online.com





Calpain 6 Protein (CAPN6) (Myc-DYKDDDDK Tag)



Image



| Overview | |
|-------------------------------|---|
| Quantity: | 20 μg |
| Target: | Calpain 6 (CAPN6) |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This Calpain 6 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |
| Product Details | |
| Characteristics: | Recombinant human Calpain-6 protein expressed in HEK293 cells. Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| Target Details | |
| Target: | Calpain 6 (CAPN6) |
| Alternative Name: | Calpain-6 (CAPN6 Products) |
| Background: | Calpains are ubiquitous, well-conserved family of calcium-dependent, cysteine proteases. The |
| | calpain proteins are heterodimers consisting of an invariant small subunit and variable large |
| | subunits. The large subunit possesses a cysteine protease domain, and both subunits possess |
| | calcium-binding domains. Calpains have been implicated in neurodegenerative processes, as |
| | their activation can be triggered by calcium influx and oxidative stress. The protein encoded by |

Target Details

| this gene is highly expressed in the placenta. Its C-terminal region lacks any homology to the |
|--|
| calmodulin-like domain of other calpains. The protein lacks critical active site residues and thus |
| is suggested to be proteolytically inactive. The protein may play a role in tumor formation by |
| inhibiting apoptosis and promoting angiogenesis. |

| Molecular Weight: | 74.4 kDa |
|-------------------|----------|
| Molecular Weight: | 74.4 kDa |

NCBI Accession: NP_055104


Application Details

| Application Notes: | Recombinant human proteins can be used for: |
|--------------------|--|
| | Native antigens for optimized antibody production |
| | Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |

Handling

| Concentration: | 50 μg/mL |
|------------------|---|
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

Images



Western Blotting

Image 1. Validation with Western Blot