antibodies

Datasheet for ABIN2713268 CBY1/PGEA1 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)

Image



Overview

Protein Characteristics:Transcript Variant 1Origin:HumanSource:HEK-293 CellsProtein Type:RecombinantPurification tag / Conjugate:This CBY1/PGEA1 protein is labelled with Myc-DYKDDDDK Tag.Application:Antibody Production (AbP), Standard (STD)Product Details.Characteristics:. Recombinant human CBY1 (transcript variant 1) protein expressed in HEK293 cells. . Produced with end-sequenced ORF clonePurity:>80 % as determined by SDS-PAGE and Coomassie blue stainingTarget Details		
Protein Characteristics:Transcript Variant 1Origin:HumanSource:HEK-293 CellsProtein Type:RecombinantPurification tag / Conjugate:This CBY1/PGEA1 protein is labelled with Myc-DYKDDDDK Tag.Application:Antibody Production (AbP), Standard (STD)Product Details.Characteristics:. Recombinant human CBY1 (transcript variant 1) protein expressed in HEK293 cells. . Produced with end-sequenced ORF clonePurity:> 80 % as determined by SDS-PAGE and Coomassie blue stainingTarget Details.Target:CBY1/PGEA1 (CBY1)Alternative Name:Cby1 (CBY1 Products)Background:Beta-catenin is a transcriptional activator and oncoprotein involved in the development of several cancers. The protein encoded by this gene interacts directly with the C-terminal region of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by	Quantity:	20 µg
Origin:HumanSource:HEK-293 CellsProtein Type:RecombinantPurification tag / Conjugate:This CBY1/PGEA1 protein is labelled with Myc-DYKDDDDK Tag.Application:Antibody Production (AbP), Standard (STD)Product DetailsProduct DetailsCharacteristics:. Recombinant human CBY1 (transcript variant 1) protein expressed in HEK293 cells. . Produced with end-sequenced ORF clonePurity:>80 % as determined by SDS-PAGE and Coomassie blue stainingTarget:CBY1/PGEA1 (CBY1)Atternative Name:CDy1 (CBY1 Products)Background:Beta-catenin is a transcriptional activator and oncoprotein involved in the development of several cancers. The protein encoded by this gene interacts directly with the C-terminal region of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by	Target:	CBY1/PGEA1 (CBY1)
Source:HEK-293 CellsProtein Type:RecombinantPurification tag / Conjugate:This CBY1/PGEA1 protein is labelled with Myc-DYKDDDDK Tag.Application:Antibody Production (AbP), Standard (STD)Product Details.Characteristics:. Recombinant human CBY1 (transcript variant 1) protein expressed in HEK293 cells. . Produced with end-sequenced ORF clonePurity:.80 % as determined by SDS-PAGE and Coomassie blue stainingTarget Details.Target:CBY1/PGEA1 (CBY1)Alternative Name:Cby1 (CBY1 Products)Background:Beta-catenin is a transcriptional activator and oncoprotein involved in the development of several cancers. The protein encoded by this gene interacts directly with the C-terminal region of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by	Protein Characteristics:	Transcript Variant 1
Protein Type:RecombinantPurification tag / Conjugate:This CBY1/PGEA1 protein is labelled with Myc-DYKDDDDK Tag.Application:Antibody Production (AbP), Standard (STD)Product Details-Characteristics:: Recombinant human CBY1 (transcript variant 1) protein expressed in HEK293 cells. · Produced with end-sequenced ORF clonePurity:> 80 % as determined by SDS-PAGE and Coomassie blue stainingTarget Details-Target:CBY1/PGEA1 (CBY1)Alternative Name:Cby1 (CBY1 Products)Background:Beta-catenin is a transcriptional activator and oncoprotein involved in the development of several cancers. The protein encoded by this gene interacts directly with the C-terminal region of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by	Origin:	Human
Purification tag / Conjugate:This CBY1/PGEA1 protein is labelled with Myc-DYKDDDDK Tag.Application:Antibody Production (AbP), Standard (STD)Product DetailsCharacteristics:· Recombinant human CBY1 (transcript variant 1) protein expressed in HEK293 cells. · Produced with end-sequenced ORF clonePurity:> 80 % as determined by SDS-PAGE and Coomassie blue stainingTarget DetailsTarget:CBY1/PGEA1 (CBY1)Alternative Name:Cby1 (CBY1 Products)Background:Beta-catenin is a transcriptional activator and oncoprotein involved in the development of several cancers. The protein encoded by this gene interacts directly with the C-terminal region of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by	Source:	HEK-293 Cells
Application:Antibody Production (AbP), Standard (STD)Product DetailsCharacteristics:· Recombinant human CBY1 (transcript variant 1) protein expressed in HEK293 cells. · Produced with end-sequenced ORF clonePurity:> 80 % as determined by SDS-PAGE and Coomassie blue stainingTarget DetailsTarget:CBY1/PGEA1 (CBY1)Alternative Name:Cby1 (CBY1 Products)Background:Beta-catenin is a transcriptional activator and oncoprotein involved in the development of several cancers. The protein encoded by this gene interacts directly with the C-terminal region of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by	Protein Type:	Recombinant
Product DetailsCharacteristics:. Recombinant human CBY1 (transcript variant 1) protein expressed in HEK293 cells. . Produced with end-sequenced ORF clonePurity:> 80 % as determined by SDS-PAGE and Coomassie blue stainingTarget DetailsTarget:CBY1/PGEA1 (CBY1)Alternative Name:Cby1 (CBY1 Products)Background:Beta-catenin is a transcriptional activator and oncoprotein involved in the development of several cancers. The protein encoded by this gene interacts directly with the C-terminal region of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by	Purification tag / Conjugate:	This CBY1/PGEA1 protein is labelled with Myc-DYKDDDDK Tag.
Characteristics: Recombinant human CBY1 (transcript variant 1) protein expressed in HEK293 cells.Produced with end-sequenced ORF clone Purity: > 80 % as determined by SDS-PAGE and Coomassie blue staining Target Details Target:CBY1/PGEA1 (CBY1) Alternative Name:Cby1 (CBY1 Products) Background:Beta-catenin is a transcriptional activator and oncoprotein involved in the development of several cancers. The protein encoded by this gene interacts directly with the C-terminal region of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by	Application:	Antibody Production (AbP), Standard (STD)
Characteristics: Recombinant human CBY1 (transcript variant 1) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone Purity: 80 % as determined by SDS-PAGE and Coomassie blue staining Target Details Target: CBY1/PGEA1 (CBY1) Alternative Name: Cby1 (CBY1 Products) Beta-catenin is a transcriptional activator and oncoprotein involved in the development of several cancers. The protein encoded by this gene interacts directly with the C-terminal region of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by 	Product Details	
Produced with end-sequenced ORF clonePurity:> 80 % as determined by SDS-PAGE and Coomassie blue stainingTarget DetailsTarget:CBY1/PGEA1 (CBY1)Alternative Name:Cby1 (CBY1 Products)Background:Beta-catenin is a transcriptional activator and oncoprotein involved in the development of several cancers. The protein encoded by this gene interacts directly with the C-terminal region of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by		
Target DetailsTarget:CBY1/PGEA1 (CBY1)Alternative Name:Cby1 (CBY1 Products)Background:Beta-catenin is a transcriptional activator and oncoprotein involved in the development of several cancers. The protein encoded by this gene interacts directly with the C-terminal region of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by	Characteristics:	
Target:CBY1/PGEA1 (CBY1)Alternative Name:Cby1 (CBY1 Products)Background:Beta-catenin is a transcriptional activator and oncoprotein involved in the development of several cancers. The protein encoded by this gene interacts directly with the C-terminal region of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by	Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Alternative Name: Cby1 (CBY1 Products) Background: Beta-catenin is a transcriptional activator and oncoprotein involved in the development of several cancers. The protein encoded by this gene interacts directly with the C-terminal region of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by	Target Details	
Background: Beta-catenin is a transcriptional activator and oncoprotein involved in the development of several cancers. The protein encoded by this gene interacts directly with the C-terminal region of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by	Target:	CBY1/PGEA1 (CBY1)
several cancers. The protein encoded by this gene interacts directly with the C-terminal region of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by	Alternative Name:	Cby1 (CBY1 Products)
of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by	Background:	Beta-catenin is a transcriptional activator and oncoprotein involved in the development of
		several cancers. The protein encoded by this gene interacts directly with the C-terminal region
		of beta-catenin, inhibiting oncogenic beta-catenin-mediated transcriptional activation by


Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN2713268 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

	encoding different isoforms have been found for this gene.
Molecular Weight:	14.3 kDa
NCBI Accession:	NP_056188
Application Details	
Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays
Comment:	
Comment: Restrictions:	Positive controls in ELISA and other antibody assays

Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze
	immediately. Only 2-3 freeze thaw cycles are recommended.

Images



Western Blotting

Image 1. Validation with Western Blot

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN2713268 | 09/11/2023 | Copyright antibodies-online. All rights reserved.