

## Datasheet for ABIN2713452

## **DDR1 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)**





| Go | to | Produ | ıct   | nac | ı |
|----|----|-------|-------|-----|---|
| OO | ιO | riout | a C L | pay |   |


| Overview                      |   |  |  |
|-------------------------------|---|--|--|
| Quantity:                     | 20 μg   |  |  |
| Target:                       | DDR1  |  |  |
| Protein Characteristics:      | Transcript Variant 1  |  |  |
| Origin:                       | Human   |  |  |
| Source:                       | HEK-293 Cells   |  |  |
| Protein Type:                 | Recombinant   |  |  |
| Purification tag / Conjugate: | This DDR1 protein is labelled with Myc-DYKDDDDK Tag.  |  |  |
| Application:                  | Antibody Production (AbP), Standard (STD)   |  |  |
| Product Details               |   |  |  |
| Characteristics:              | <ul> <li>Recombinant human CD167a / DDR1 (transcript variant 1) protein expressed in HEK293 cells.</li> </ul> |  |  |
|                               | Produced with end-sequenced ORF clone   |  |  |
| Purity:                       | > 80 % as determined by SDS-PAGE and Coomassie blue staining  |  |  |
| Target Details                |   |  |  |
| Target:                       | DDR1  |  |  |
| Alternative Name:             | Cd167a,ddr1 (DDR1 Products)   |  |  |
| Background:                   | Receptor tyrosine kinases play a key role in the communication of cells with their                            |  |  |
|                               | microenvironment. These kinases are involved in the regulation of cell growth, differentiation                |  |  |
|                               | and metabolism. The protein encoded by this gene belongs to a subfamily of tyrosine kinase                    |  |  |

Storage Comment:

|                     | receptors with homology to Dictyostelium discoideum protein discoidin I in their extracellular domain, and that are activated by various types of collagen. Expression of this protein is restricted to epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In addition, it has been shown to be significantly overexpressed in several human tumors. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. |  |
|---------------------|---|--|
| Molecular Weight:   | 100.9 kDa   |  |
| NCBI Accession:     | NP_054699   |  |
| Pathways:           | RTK Signaling, Smooth Muscle Cell Migration   |  |
| Application Details |   |  |
| Application Notes:  | Recombinant human proteins can be used for:   |  |
|                     | Native antigens for optimized antibody production   |  |
|                     | Positive controls in ELISA and other antibody assays  |  |
| Comment:            | The tag is located at the C-terminal.   |  |
| Restrictions:       | For Research Use only   |  |
| Handling            |   |  |
| Concentration:      | 50 μg/mL  |  |
| Buffer:             | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.  |  |
| Storage:            | age: -80 °C   |  |
|                     |   |  |

immediately. Only 2-3 freeze thaw cycles are recommended.

Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze



## **Western Blotting**

Image 1. Validation with Western Blot