

Datasheet for ABIN2713665

FGFR2 Protein (DYKDDDDK-His Tag)





| \sim | | | |
|--------|-----|-------|----|
| | ve | r\/ | ٨ |
| \cup | V C | 1 V I | ٧V |

| 20 μg | |
|--|--|
| FGFR2 | |
| Human | |
| HEK-293 Cells | |
| Recombinant | |
| This FGFR2 protein is labelled with DYKDDDDK-His Tag. | |
| Antibody Production (AbP), Standard (STD) | |
| | |
| Recombinant human CD332 / FGFR-2 (C-term DDK/His tag, transcript variant 1) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone | |
| > 80 % as determined by SDS-PAGE and Coomassie blue staining | |
| | |
| FGFR2 | |
| Cd332,fgfr-2 (FGFR2 Products) | |
| The protein encoded by this gene is a member of the fibroblast growth factor receptor family where amino acid sequence is highly conserved between members and throughout evolution FGFR family members differ from one another in their ligand affinities and tissue distribution full-length representative protein consists of an extracellular region, composed of three | |
| | |

| immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a |
|---|
| cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with |
| fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately |
| influencing mitogenesis and differentiation. This particular family member is a high-affinity |
| receptor for acidic, basic and/or keratinocyte growth factor, depending on the isoform. |
| Mutations in this gene are associated with Crouzon syndrome, Pfeiffer syndrome, |
| Craniosynostosis, Apert syndrome, Jackson-Weiss syndrome, Beare-Stevenson cutis gyrata |
| syndrome, Saethre-Chotzen syndrome, and syndromic craniosynostosis. Multiple alternatively |
| spliced transcript variants encoding different isoforms have been noted for this gene. |
| |


| Molecular Weight: | 42 kDa |
|-------------------|--|
| NCBI Accession: | NP_000132 |
| Pathways: | RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber |
| | Development, Growth Factor Binding |

Application Details

| Application Notes: | Recombinant human proteins can be used for: | |
|--------------------|--|--|
| | Native antigens for optimized antibody production | |
| | Positive controls in ELISA and other antibody assays | |
| Comment: | The tag is located at the C-terminal. | |
| Restrictions: | For Research Use only | |

Handling

| Concentration: | 50 μg/mL | |
|------------------|--|--|
| Buffer: | 1 x PBS, pH 7.4, 10 % glycerol. | |
| Storage: | -80 °C | |
| Storage Comment: | age Comment: Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. | |



Western Blotting

Image 1. Validation with Western Blot