# antibodies -online.com





## **CEACAM3 Protein (Myc-DYKDDDDK Tag)**



Image



Go to Product page

$\sim$				
	$ V \cap$	r\/I	19	٨

molecules (CEACAMs), which are used by several bacterial pathogens to bind and cells. The encoded transmembrane protein directs phagocytosis of several bacte		
Origin: Human  Source: HEK-293 Cells  Protein Type: Recombinant  Purification tag / Conjugate: This CEACAM3 protein is labelled with Myc-DYKDDDDK Tag.  Application: Antibody Production (AbP), Standard (STD)  Product Details  Characteristics: • Recombinant human CD66d / CEACAM3 protein expressed in HEK293 cells. • Produced with end-sequenced ORF clone  Purity: > 80 % as determined by SDS-PAGE and Coomassie blue staining  Target Details  Target: CEACAM3  Alternative Name: Cd66d,ceacam3 (CEACAM3 Products)  Background: This gene encodes a member of the family of carcinoembryonic antigen-related of molecules (CEACAMs), which are used by several bacterial pathogens to bind and cells. The encoded transmembrane protein directs phagocytosis of several bacterial pathogens.	uantity:	
Source: HEK-293 Cells  Protein Type: Recombinant  Purification tag / Conjugate: This CEACAM3 protein is labelled with Myc-DYKDDDDK Tag.  Application: Antibody Production (AbP), Standard (STD)  Product Details  Characteristics: • Recombinant human CD66d / CEACAM3 protein expressed in HEK293 cells. • Produced with end-sequenced ORF clone  Purity: > 80 % as determined by SDS-PAGE and Coomassie blue staining  Target Details  Target: CEACAM3  Alternative Name: Cd66d,ceacam3 (CEACAM3 Products)  Background: This gene encodes a member of the family of carcinoembryonic antigen-related of molecules (CEACAMs), which are used by several bacterial pathogens to bind and cells. The encoded transmembrane protein directs phagocytosis of several bacter.	arget:	CEACAM3
Protein Type: Recombinant  Purification tag / Conjugate: This CEACAM3 protein is labelled with Myc-DYKDDDDK Tag.  Application: Antibody Production (AbP), Standard (STD)  Product Details  Characteristics: • Recombinant human CD66d / CEACAM3 protein expressed in HEK293 cells. • Produced with end-sequenced ORF clone  Purity: > 80 % as determined by SDS-PAGE and Coomassie blue staining  Target Details  Target: CEACAM3  Alternative Name: Cd66d,ceacam3 (CEACAM3 Products)  Background: This gene encodes a member of the family of carcinoembryonic antigen-related of molecules (CEACAMs), which are used by several bacterial pathogens to bind and cells. The encoded transmembrane protein directs phagocytosis of several bacterial	igin:	Human
Purification tag / Conjugate: This CEACAM3 protein is labelled with Myc-DYKDDDDK Tag.  Application: Antibody Production (AbP), Standard (STD)  Product Details  Characteristics: • Recombinant human CD66d / CEACAM3 protein expressed in HEK293 cells. • Produced with end-sequenced ORF clone  Purity: > 80 % as determined by SDS-PAGE and Coomassie blue staining  Target Details  Target: CEACAM3  Alternative Name: Cd66d,ceacam3 (CEACAM3 Products)  Background: This gene encodes a member of the family of carcinoembryonic antigen-related of molecules (CEACAMs), which are used by several bacterial pathogens to bind and cells. The encoded transmembrane protein directs phagocytosis of several bacterial pathogens to several bacterial pathogens.	ource:	HEK-293 Cells
Application: Antibody Production (AbP), Standard (STD)  Product Details  Characteristics: Produced with human CD66d / CEACAM3 protein expressed in HEK293 cells. Produced with end-sequenced ORF clone  Purity: > 80 % as determined by SDS-PAGE and Coomassie blue staining  Target Details  Target: CEACAM3  Alternative Name: Cd66d,ceacam3 (CEACAM3 Products)  Background: This gene encodes a member of the family of carcinoembryonic antigen-related of molecules (CEACAMs), which are used by several bacterial pathogens to bind and cells. The encoded transmembrane protein directs phagocytosis of several bacter	otein Type:	Recombinant
Product Details  - Recombinant human CD66d / CEACAM3 protein expressed in HEK293 cells Produced with end-sequenced ORF clone  - Purity: - > 80 % as determined by SDS-PAGE and Coomassie blue staining  - Target Details  - Target: - CEACAM3  - Alternative Name: - Cd66d,ceacam3 (CEACAM3 Products)  - Background: - This gene encodes a member of the family of carcinoembryonic antigen-related of molecules (CEACAMs), which are used by several bacterial pathogens to bind and cells. The encoded transmembrane protein directs phagocytosis of several bacterial pathogens.	ırification tag / Conjugate:	This CEACAM3 protein is labelled with Myc-DYKDDDDK Tag.
Characteristics:  • Recombinant human CD66d / CEACAM3 protein expressed in HEK293 cells. • Produced with end-sequenced ORF clone  Purity:  > 80 % as determined by SDS-PAGE and Coomassie blue staining  Target Details  Target:  CEACAM3  Alternative Name:  Cd66d,ceacam3 (CEACAM3 Products)  Background:  This gene encodes a member of the family of carcinoembryonic antigen-related of molecules (CEACAMs), which are used by several bacterial pathogens to bind and cells. The encoded transmembrane protein directs phagocytosis of several bacterial pathogens.	pplication:	Antibody Production (AbP), Standard (STD)
Purity: > 80 % as determined by SDS-PAGE and Coomassie blue staining  Target Details  Target: CEACAM3  Alternative Name: Cd66d,ceacam3 (CEACAM3 Products)  Background: This gene encodes a member of the family of carcinoembryonic antigen-related complexules (CEACAMs), which are used by several bacterial pathogens to bind and cells. The encoded transmembrane protein directs phagocytosis of several bacterial pathogens.	roduct Details	
Target Details  Target: CEACAM3  Alternative Name: Cd66d,ceacam3 (CEACAM3 Products)  Background: This gene encodes a member of the family of carcinoembryonic antigen-related of molecules (CEACAMs), which are used by several bacterial pathogens to bind and cells. The encoded transmembrane protein directs phagocytosis of several bacterial pathogens.		
Target: CEACAM3  Alternative Name: Cd66d,ceacam3 (CEACAM3 Products)  Background: This gene encodes a member of the family of carcinoembryonic antigen-related of molecules (CEACAMs), which are used by several bacterial pathogens to bind and cells. The encoded transmembrane protein directs phagocytosis of several bacterial pathogens.	ırity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Alternative Name: Cd66d,ceacam3 (CEACAM3 Products)  Background: This gene encodes a member of the family of carcinoembryonic antigen-related of molecules (CEACAMs), which are used by several bacterial pathogens to bind and cells. The encoded transmembrane protein directs phagocytosis of several bacterial pathogens.	arget Details	
Background:  This gene encodes a member of the family of carcinoembryonic antigen-related of molecules (CEACAMs), which are used by several bacterial pathogens to bind and cells. The encoded transmembrane protein directs phagocytosis of several bacterial pathogens.	arget:	CEACAM3
molecules (CEACAMs), which are used by several bacterial pathogens to bind and cells. The encoded transmembrane protein directs phagocytosis of several bacte	ternative Name:	Cd66d,ceacam3 (CEACAM3 Products)
		This gene encodes a member of the family of carcinoembryonic antigen-related cell adhesion molecules (CEACAMs), which are used by several bacterial pathogens to bind and invade host cells. The encoded transmembrane protein directs phagocytosis of several bacterial species that is dependent on the small GTPase Rac. It is thought to serve an important role in controlling human-specific pathogens by the innate immune system. Alternatively spliced

### Target Details

	transcript variants have been described.
Molecular Weight:	23.4 kDa
NCBI Accession:	NP_001806


## **Application Details**

Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only

### Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.

#### **Images**



#### **Western Blotting**

Image 1. Validation with Western Blot