

Datasheet for ABIN2713999

## CDYL Protein (Transcript Variant 3) (Myc-DYKDDDDK Tag)



[Go to Product page](#)

### 1 Image

#### Overview

|                               |  |
|-------------------------------|--|
| Quantity:                     | 20 µg  |
| Target:                       | CDYL   |
| Protein Characteristics:      | Transcript Variant 3                                 |
| Origin:                       | Human  |
| Source:                       | HEK-293 Cells  |
| Protein Type:                 | Recombinant  |
| Purification tag / Conjugate: | This CDYL protein is labelled with Myc-DYKDDDDK Tag. |
| Application:                  | Antibody Production (AbP), Standard (STD)            |

#### Product Details

|                  |  |
|------------------|--|
| Characteristics: | <ul style="list-style-type: none"> <li>• Recombinant human Chromodomain protein, Y-like (CDYL), transcript variant 3 (transcript variant 3) protein expressed in HEK293 cells.</li> <li>• Produced with end-sequenced ORF clone</li> </ul> |
| Purity:          | > 80 % as determined by SDS-PAGE and Coomassie blue staining   |

#### Target Details

|             |   |
|-------------|---|
| Target:     | CDYL  |
| Abstract:   | <a href="#">CDYL Products</a>   |
| Background: | Chromodomain Y is a primate-specific Y-chromosomal gene family expressed exclusively in the testis and implicated in infertility. Although the Y-linked genes are testis-specific, this autosomal gene is ubiquitously expressed. The Y-linked genes arose by retrotransposition of an mRNA |

## Target Details

from this gene, followed by amplification of the retroposed gene. Proteins encoded by this gene superfamily possess a chromodomain, a motif implicated in chromatin binding and gene suppression, and a catalytic domain believed to be involved in histone acetylation. Multiple proteins are encoded by transcript variants of this gene.

Molecular Weight: 34.4 kDa

NCBI Accession: [NP\\_736608](#)

## Application Details

Application Notes: Recombinant human proteins can be used for:  
Native antigens for optimized antibody production  
Positive controls in ELISA and other antibody assays

Comment: The tag is located at the C-terminal.

Restrictions: For Research Use only

## Handling


Concentration: 50 µg/mL

Buffer: 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.

Storage: -80 °C

Storage Comment: Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.

## Images



### Western Blotting

**Image 1.** Validation with Western Blot