# antibodies .- online.com





# ABHD2 Protein (Transcript Variant 2) (Myc-DYKDDDDK Tag)



Overview

Image



Go to Product page

| Quantity:                | 20 μg                |
|--------------------------|----------------------|
| Target:                  | ABHD2                |
| Protein Characteristics: | Transcript Variant 2 |
| Origin:                  | Human                |
| Source:                  | HEK-293 Cells        |
|                          |                      |

| Purification tag / Conjugate: | This ABHD2 protein is labelled with Myc-DYKDDDDK Tag. |
|-------------------------------|---|

Application: Antibody Production (AbP), Standard (STD)

Recombinant

#### **Product Details**

Protein Type:

| Characteristics: | <ul> <li>Recombinant human ABHD2 / LABH2 (transcript variant 2) protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul> |
|------------------|--|
| Purity:          | > 80 % as determined by SDS-PAGE and Coomassie blue staining   |

#### **Target Details**

| Target:           | ABHD2   |
|-------------------|---|
| Alternative Name: | Abhd2,labh2 (ABHD2 Products)  |
| Background:       | This gene encodes a protein containing an alpha/beta hydrolase fold, which is a catalytic domain found in a wide range of enzymes. The encoded protein is an acylglycerol lipase that |
|                   | catalyzes the hydrolysis of endocannabinoid arachidonoylglycerol from the cell membrane.  |

# **Target Details**

|                   | This leads to activation of the sperm calcium channel CatSper, which results in sperm              |
|-------------------|--|
|                   | activation. Alternative splicing of this gene results in two transcript variants encoding the same |
|                   | protein.   |
| Molecular Weight: | 48.1 kDa   |
| NCBI Accession:   | NP 690888  |


# **Application Details**

| Application Notes: | Recombinant human proteins can be used for:          |
|--------------------|--|
|                    | Native antigens for optimized antibody production    |
|                    | Positive controls in ELISA and other antibody assays |
| Comment:           | The tag is located at the C-terminal.                |
| Restrictions:      | For Research Use only                                |

# Handling

| Concentration:   | 50 μg/mL  |
|------------------|---|
| Buffer:          | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.  |
| Storage:         | -80 °C  |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

## **Images**



## **Western Blotting**

Image 1. Validation with Western Blot