

Datasheet for ABIN2714690

ANGPTL3 Protein (His tag)






Overview

Quantity:	50 µg
Target:	ANGPTL3
Origin:	Human
Source:	CHO Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This ANGPTL3 protein is labelled with His tag.
Application:	Antibody Production (AbP), Standard (STD), Functional Studies (Func), Protein Interaction (PI)
Product Details	
Specificity:	Optimal preservation of protein structure, post-translational modifications and functions.
Specificity: Characteristics:	Optimal preservation of protein structure, post-translational modifications and functions. Recombinant human Angiopoietin-5 protein expressed in CHO cells.
· · · · · · · · · · · · · · · · · · ·	Recombinant human Angiopoietin-5 protein expressed in CHO cells.
· · · · · · · · · · · · · · · · · · ·	 Recombinant human Angiopoietin-5 protein expressed in CHO cells. Produced with end-sequenced ORF clone
Characteristics:	 Recombinant human Angiopoietin-5 protein expressed in CHO cells. Produced with end-sequenced ORF clone Tested for bioactivity.
Characteristics: Purity:	 Recombinant human Angiopoietin-5 protein expressed in CHO cells. Produced with end-sequenced ORF clone Tested for bioactivity. > 95 % as determined by SDS-PAGE and Coomassie blue staining
Characteristics: Purity: Endotoxin Level:	 Recombinant human Angiopoietin-5 protein expressed in CHO cells. Produced with end-sequenced ORF clone Tested for bioactivity. > 95 % as determined by SDS-PAGE and Coomassie blue staining Endotoxin level is <0.1 ng/μg of protein (<1EU/μg).
Characteristics: Purity: Endotoxin Level: Biological Activity Comment:	 Recombinant human Angiopoietin-5 protein expressed in CHO cells. Produced with end-sequenced ORF clone Tested for bioactivity. > 95 % as determined by SDS-PAGE and Coomassie blue staining Endotoxin level is <0.1 ng/μg of protein (<1EU/μg).

Target Details

rarget betano	
Alternative Name:	Angiopoietin-5 (ANGPTL3 Products)
Background:	This gene encodes a member of a family of secreted proteins that function in angiogenesis.
	The encoded protein, which is expressed predominantly in the liver, is further processed into ar
	N-terminal coiled-coil domain-containing chain and a C-terminal fibrinogen chain. The N-
	terminal chain is important for lipid metabolism, while the C-terminal chain may be involved in
	angiogenesis. Mutations in this gene cause familial hypobetalipoproteinemia type 2.
Molecular Weight:	53.64
NCBI Accession:	NP_055310
Application Details	
Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
	Protein-protein interaction
	In vitro biochemical assays and cell-based functional assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only
Handling	
Buffer:	Lyophilized from a 0.2 μM filtered solution of 20 mM phosphate buffer,100 mM NaCl, pH 7.2
Handling Advice:	Resuspend the protein in the desired concentration in proper buffer
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze
	immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot