

Datasheet for ABIN2714747

Annexin VII Protein (Transcript Variant 2) (Myc-DYKDDDDK Tag)



Go to Product pag

1 Image

Overview	
Quantity:	20 μg
Target:	Annexin VII (ANXA7)
Protein Characteristics:	Transcript Variant 2
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Annexin VII protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	Recombinant human Annexin A7 / ANXA7 (transcript variant 2) protein expressed in HEK293
	cells. • Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	Annexin VII (ANXA7)
Alternative Name:	Annexin a7,anxa7 (ANXA7 Products)
Background:	Annexin VII is a member of the annexin family of calcium-dependent phospholipid binding
	proteins. The Annexin VII gene contains 14 exons and spans approximately 34 kb of DNA. An
	alternatively spliced cassette exon results in two mRNA transcripts of 2.0 and 2.4 kb which are

predicted to generate two protein isoforms differing in their N-terminal domain. The alternative splicing event is tissue specific and the mRNA containing the cassette exon is prevalent in brain, heart and skeletal muscle. The transcripts also differ in their 3'-non coding regions by the use of two alternative poly(A) signals. Annexin VII encodes a protein with a molecular weight of approximately 51 kDa with a unique, highly hydrophobic N-terminal domain of 167 amino acids and a conserved C-terminal region of 299 amino acids. The latter domain is composed of alternating hydrophobic and hydrophilic segments. Structural analysis of the protein suggests that Annexin VII is a membrane binding protein with diverse properties, including voltage-sensitive calcium channel activity, ion selectivity and membrane fusion.

Molecular Weight:

52.6 kDa

NCBI Accession:

NP_004025

Application Details

A 1.		A 1 .
/\nnli	nation	Notes:
	Jaulon	INULES.

Recombinant human proteins can be used for:

Native antigens for optimized antibody production

Positive controls in ELISA and other antibody assays

Comment:

The tag is located at the C-terminal.

Restrictions:

For Research Use only

Handling

Concentr	ation:

50 μg/mL

Buffer:

25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.


Storage:

-80 °C

Storage Comment:

Store at -80 $^{\circ}\text{C}.$ Thaw on ice, aliquot to individual single-use tubes, and then re-freeze

immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot