

Datasheet for ABIN2714759

AMH Protein (Myc-DYKDDDDK Tag)**1** Image**1** Publication[Go to Product page](#)

Overview

Quantity:	20 µg
Target:	AMH
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This AMH protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)

Product Details

Characteristics:	<ul style="list-style-type: none">• Recombinant human Anti-Muellerian Hormone / AMH protein expressed in HEK293 cells.• Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining

Target Details

Target:	AMH
Alternative Name:	Anti-Muellerian Hormone, amh (AMH Products)
Background:	<p>This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression.</p> <p>The encoded preproprotein is proteolytically processed to generate N- and C-terminal cleavage products that homodimerize and associate to form a biologically active noncovalent complex.</p>

Target Details

This complex binds to the anti-Mullerian hormone receptor type 2 and causes the regression of Mullerian ducts in the male embryo that would otherwise differentiate into the uterus and fallopian tubes. This protein also plays a role in Leydig cell differentiation and function and follicular development in adult females. Mutations in this gene result in persistent Mullerian duct syndrome.

Molecular Weight: 56.6 kDa

NCBI Accession: [NP_000470](#)

Pathways: [Negative Regulation of Hormone Secretion](#)

Application Details

Application Notes: Recombinant human proteins can be used for:
Native antigens for optimized antibody production
Positive controls in ELISA and other antibody assays

Comment: The tag is located at the C-terminal.

Restrictions: For Research Use only

Handling

Concentration: 50 µg/mL

Buffer: 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.

Storage: -80 °C

Storage Comment: Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.

Publications

Product cited in: Shiryayev, Aleshin, Muranaka, Kukreja, Routenberg, Remacle, Liddington, Cieplak, Kozlov, Strongin: "Structural and functional diversity of metalloproteinases encoded by the Bacteroides fragilis pathogenicity island." in: **The FEBS journal**, Vol. 281, Issue 11, pp. 2487-502, (2014) ([PubMed](#)).


Ge, Siegel, Jordan, Naumann: "Ligand binding alters dimerization and sequestering of urokinase receptors in raft-mimicking lipid mixtures." in: **Biophysical journal**, Vol. 107, Issue 9, pp. 2101-11, (2014) ([PubMed](#)).

Garcia-Sanz, Quintanilla, Lafita, Moreno-Bueno, García-Gutierrez, Tabor, Varela, Shiio, Larsson, Portillo, Leon: "Sin3b interacts with Myc and decreases Myc levels." in: **The Journal of biological chemistry**, Vol. 289, Issue 32, pp. 22221-36, (2014) ([PubMed](#)).

Wang, Henry, Distefano, Wang, Räikkönen, Mönkkönen, Tanaka, Morita: "Butyrophilin 3A1 plays an essential role in prenyl pyrophosphate stimulation of human V α 2V β 2 T cells." in: **Journal of immunology (Baltimore, Md. : 1950)**, Vol. 191, Issue 3, pp. 1029-42, (2013) ([PubMed](#)).

Bardeleben, Sharma, Reeve, Bassilian, Frost, Hoang, Shi, Lichtenstein: "Metabolomics identifies pyrimidine starvation as the mechanism of 5-aminoimidazole-4-carboxamide-1- β -ribose-induced apoptosis in multiple myeloma cells." in: **Molecular cancer therapeutics**, Vol. 12, Issue 7, pp. 1310-21, (2013) ([PubMed](#)).

Images



Western Blotting

Image 1. Validation with Western Blot