antibodies - online.com





APOBEC3C Protein (Myc-DYKDDDDK Tag)





| Go to | Prod | uct | page |
|-------|------|-----|------|
| | | | |

| \sim | | | | |
|--------|----|-----------|----|-----|
| | VE | ۲۱د | // | ١٨. |
| \cup | V | ر ار ا | ۷ľ | Vν |

| Overview | |
|-------------------------------|---|
| Quantity: | 20 μg |
| Target: | APOBEC3C |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This APOBEC3C protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |
| Product Details | |
| Characteristics: | Recombinant human APOBEC3C protein expressed in HEK293 cells. Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| Target Details | |
| Target: | APOBEC3C |
| Alternative Name: | Apobec3c (APOBEC3C Products) |
| Background: | This gene is a member of the cytidine deaminase gene family. It is one of seven related genes or pseudogenes found in a cluster thought to result from gene duplication, on chromosome 22. Members of the cluster encode proteins that are structurally and functionally related to the C to U RNA-editing cytidine deaminase APOBEC1. It is thought that the proteins may be RNA editing enzymes and have roles in growth or cell cycle control. |

Target Details

| Molecular Weight: | 22.6 kDa |
|-------------------|-----------|
| NCBI Accession: | NP_055323 |


Application Details

| Application Notes: | Recombinant human proteins can be used for: |
|--------------------|--|
| | Native antigens for optimized antibody production |
| | Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |

Handling

| Concentration: | 50 μg/mL |
|------------------|---|
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

Images



Western Blotting

Image 1. Validation with Western Blot