# antibodies -online.com





## **Arx Protein (Myc-DYKDDDDK Tag)**



Image



Go to Product page

| $\sim$ |           |      |    |   |
|--------|-----------|------|----|---|
|        | $ V \cap$ | r\/I | 19 | ٨ |

| Overview                      |   |
|-------------------------------|---|
| Quantity:                     | 20 μg   |
| Target:                       | Arx (ARX)   |
| Origin:                       | Human   |
| Source:                       | HEK-293 Cells   |
| Protein Type:                 | Recombinant   |
| Purification tag / Conjugate: | This Arx protein is labelled with Myc-DYKDDDDK Tag.   |
| Application:                  | Antibody Production (AbP), Standard (STD)   |
| Product Details               |   |
| Characteristics:              | <ul> <li>Recombinant human ARX protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>   |
| Purity:                       | > 80 % as determined by SDS-PAGE and Coomassie blue staining  |
| Target Details                |   |
| Target:                       | Arx (ARX)   |
| Alternative Name:             | Arx (ARX Products)  |
| Background:                   | This gene is a homeobox-containing gene expressed during development. The expressed protein contains two conserved domains, a C-peptide (or aristaless domain) and the prd-like class homeobox domain. It is a member of the group-II aristaless-related protein family whose members are expressed primarily in the central and/or peripheral nervous system. This gene is thought to be involved in CNS development. Expansion of a polyalanine tract and other |

### **Target Details**

|                   | mutations in this gene cause X-linked mental retardation and epilepsy. |
|-------------------|--|
| Molecular Weight: | 58 kDa   |
| NCBI Accession:   | NP_620689  |


## **Application Details**

| Application Notes: | Recombinant human proteins can be used for:          |
|--------------------|--|
|                    | Native antigens for optimized antibody production    |
|                    | Positive controls in ELISA and other antibody assays |
| Comment:           | The tag is located at the C-terminal.                |
| Restrictions:      | For Research Use only                                |

## Handling

| Concentration:   | 50 μg/mL  |
|------------------|---|
| Buffer:          | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.  |
| Storage:         | -80 °C  |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

#### Images



#### **Western Blotting**

Image 1. Validation with Western Blot