.-online.com antibodies

Datasheet for ABIN2715013 Ataxin 1 Protein (ATXN1) (Transcript Variant 2) (Myc-DYKDDDDK Tag)






Overview

0.00.000	
Quantity:	20 µg
Target:	Ataxin 1 (ATXN1)
Protein Characteristics:	Transcript Variant 2
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Ataxin 1 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	 Recombinant human Ataxin-1 (transcript variant 2) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	Ataxin 1 (ATXN1)
Alternative Name:	Ataxin-1 (ATXN1 Products)
Background:	The autosomal dominant cerebellar ataxias (ADCA) are a heterogeneous group of
	neurodegenerative disorders characterized by progressive degeneration of the cerebellum,
	brain stem and spinal cord. Clinically, ADCA has been divided into three groups: ADCA types I-III.
	ADCAI is genetically heterogeneous, with five genetic loci, designated spinocerebellar ataxia

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN2715013 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

	(SCA) 1, 2, 3, 4 and 6, being assigned to five different chromosomes. ADCAII, which always
	presents with retinal degeneration (SCA7), and ADCAIII often referred to as the `pure' cerebellar
	syndrome (SCA5), are most likely homogeneous disorders. Several SCA genes have been
	cloned and shown to contain CAG repeats in their coding regions. ADCA is caused by the
	expansion of the CAG repeats, producing an elongated polyglutamine tract in the corresponding
	protein. The expanded repeats are variable in size and unstable, usually increasing in size when
	transmitted to successive generations. The function of the ataxins is not known. This locus has
	been mapped to chromosome 6, and it has been determined that the diseased allele contains
	40-83 CAG repeats, compared to 6-39 in the normal allele, and is associated with
	spinocerebellar ataxia type 1 (SCA1). At least two transcript variants encoding the same protein
	have been found for this gene.
Molecular Weight:	86.7 kDa
NCBI Accession:	NP_001121636
Pathways:	Synaptic Membrane
Application Details	
Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only
Handling	
Handling	
Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze
	immediately. Only 2-3 freeze thaw cycles are recommended.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN2715013 | 09/11/2023 | Copyright antibodies-online. All rights reserved.



Western Blotting

Image 1. Validation with Western Blot

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN2715013 | 09/11/2023 | Copyright antibodies-online. All rights reserved.