## antibodies -online.com





## ATP5F1D Protein (Transcript Variant 2) (Myc-DYKDDDDK Tag)



**Image** 

Background:



Go to Product page

Overview	
Quantity:	20 μg
Target:	ATP5F1D
Protein Characteristics:	Transcript Variant 2
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This ATP5F1D protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	<ul> <li>Recombinant human ATP5D (transcript variant 2) protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	ATP5F1D
Alternative Name:	Atp5d (ATP5F1D Products)


This gene encodes a subunit of mitochondrial ATP synthase. Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner

membrane during oxidative phosphorylation. ATP synthase is composed of two linked multi-

subunit complexes: the soluble catalytic core, F1, and the membrane-spanning component, Fo,

## **Target Details**

rarget Details	
	comprising the proton channel. The catalytic portion of mitochondrial ATP synthase consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled with a stoichiometry of 3 alpha, 3 beta, and a single representative of the other 3. The proton channel consists of three main subunits (a, b, c). This gene encodes the delta subunit of the catalytic core. Alternatively spliced transcript variants encoding the same isoform have been identified.
Molecular Weight:	15 kDa
NCBI Accession:	NP_001001975
Pathways:	Proton Transport, Ribonucleoside Biosynthetic Process
Application Details	
Application Notes:	Recombinant human proteins can be used for:  Native antigens for optimized antibody production  Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only
Handling	
Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.



## **Western Blotting**

Image 1. Validation with Western Blot