antibodies .- online.com





ATP5F1D Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image



Go to Product page

| \sim | |
|--------|----------|
| ()\/\ | rview |
| \cup | 1 410 44 |

| Overview | |
|-------------------------------|---|
| Quantity: | 20 μg |
| Target: | ATP5F1D |
| Protein Characteristics: | Transcript Variant 1 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This ATP5F1D protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |
| Product Details | |
| Characteristics: | Recombinant human ATP5D (transcript variant 1) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| Target Details | |
| Target: | ATP5F1D |
| Alternative Name: | Atp5d (ATP5F1D Products) |
| Background: | This gene encodes a subunit of mitochondrial ATP synthase. Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. ATP synthase is composed of two linked multi- |
| | subunit complexes: the soluble catalytic core, F1, and the membrane-spanning component, Fo, |


Target Details

Storage Comment:

| Target Details | |
|---------------------|--|
| | comprising the proton channel. The catalytic portion of mitochondrial ATP synthase consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled with a stoichiometry of 3 alpha, 3 beta, and a single representative of the other 3. The proton channel consists of three main subunits (a, b, c). This gene encodes the delta subunit of the catalytic core. Alternatively spliced transcript variants encoding the same isoform have been identified. |
| Molecular Weight: | 15 kDa |
| NCBI Accession: | NP_001678 |
| Pathways: | Proton Transport, Ribonucleoside Biosynthetic Process |
| Application Details | |
| Application Notes: | Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |
| Handling | |
| Concentration: | 50 μg/mL |
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| 0. 0 . | O |

Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze

immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot