# antibodies -online.com



Datasheet for ABIN2715050

## ATP6V0A4 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)





0 - 1	⊦ ⊸ Г	2500	110+	pad
( ¬( )	( ) F	71()(	11 16 11	nau

	rv/		

Overview		
Quantity:	20 μg	
Target:	ATP6V0A4	
Protein Characteristics:	Transcript Variant 1	
Origin:	Human	
Source:	HEK-293 Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This ATP6V0A4 protein is labelled with Myc-DYKDDDDK Tag.	
Application:	Antibody Production (AbP), Standard (STD)	
Product Details		
Characteristics:	<ul> <li>Recombinant human ATP6V0A4 (transcript variant 1) protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>	
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining	
Target Details		
Target:	ATP6V0A4	
Alternative Name:	Atp6v0a4 (ATP6V0A4 Products)	
Background:	This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of intracellular compartments of eukaryotic cells. V-ATPase dependent	
	acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-	


	ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H	
subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists o		
	different subunits: a, c, c', c", and d. This gene is one of four genes in man and mouse that	
	encode different isoforms of the a subunit. Alternatively spliced transcript variants encoding the	
	same protein have been described. Mutations in this gene are associated with renal tubular	
	acidosis associated with preserved hearing.	
Molecular Weight:	96.2 kDa	
NCBI Accession:	NP_065683	
Pathways:	Sensory Perception of Sound, Transition Metal Ion Homeostasis, Proton Transport	

### **Application Details**

Application Notes:	Recombinant human proteins can be used for:	
	Native antigens for optimized antibody production	
	Positive controls in ELISA and other antibody assays	
Comment:	The tag is located at the C-terminal.	
Restrictions:	For Research Use only	

### Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.



### **Western Blotting**

Image 1. Validation with Western Blot