

Datasheet for ABIN2715192

BCL2L1 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image




Go to Product page

Overview

Quantity:	20 μg
Target:	BCL2L1
Protein Characteristics:	Transcript Variant 1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This BCL2L1 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	 Recombinant human Bcl-2-like 1 (transcript variant 1) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone
Characteristics: Purity:	
	Produced with end-sequenced ORF clone
Purity:	Produced with end-sequenced ORF clone
Purity: Target Details	 Produced with end-sequenced ORF clone > 80 % as determined by SDS-PAGE and Coomassie blue staining
Purity: Target Details Target:	Produced with end-sequenced ORF clone > 80 % as determined by SDS-PAGE and Coomassie blue staining BCL2L1
Purity: Target Details Target: Alternative Name:	Produced with end-sequenced ORF clone > 80 % as determined by SDS-PAGE and Coomassie blue staining BCL2L1 Bcl-2-Like 1 (BCL2L1 Products)
Purity: Target Details Target: Alternative Name:	Produced with end-sequenced ORF clone > 80 % as determined by SDS-PAGE and Coomassie blue staining BCL2L1 Bcl-2-Like 1 (BCL2L1 Products) The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members form hetero- or homodimers and act as anti- or pro-apoptotic regulators that are involved in a
Purity: Target Details Target: Alternative Name:	 Produced with end-sequenced ORF clone > 80 % as determined by SDS-PAGE and Coomassie blue staining BCL2L1 Bcl-2-Like 1 (BCL2L1 Products) The protein encoded by this gene belongs to the BCL-2 protein family. BCL-2 family members

- Target Betane	
	channel (VDAC) opening. VDAC regulates mitochondrial membrane potential, and thus controls the production of reactive oxygen species and release of cytochrome C by mitochondria, both of which are the potent inducers of cell apoptosis. Alternative splicing results in multiple transcript variants encoding two different isoforms. The longer isoform acts as an apoptotic inhibitor and the shorter isoform acts as an apoptotic activator.
Molecular Weight:	25.9 kDa
NCBI Accession:	NP_612815
Pathways:	Apoptosis, Negative Regulation of intrinsic apoptotic Signaling
Application Details	
Application Notes:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only
Handling	
Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot