

Datasheet for ABIN271827

anti-STAT3 antibody**2** Images[Go to Product page](#)

Overview

Quantity:	0.1 mg
Target:	STAT3
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STAT3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Enzyme Immunoassay (EIA)

Product Details

Specificity:	This antibody detects endogenous levels of STAT3 protein. (region surrounding Thr721)
Cross-Reactivity (Details):	Species reactivity (expected):Mouse and Rat. Species reactivity (tested):Human.
Purification:	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen
Purity:	> 95 % pure by SDS-PAGE

Target Details

Target:	STAT3
Alternative Name:	STAT3 (STAT3 Products)
Background:	Membrane receptor signaling by various ligands, including interferons and growth hormones

Target Details

such as EGF, induces activation of Jak kinases which then leads to tyrosine phosphorylation of the various Stat transcription factors. Stat1 and Stat2 are induced by IFN- α and form a heterodimer which is part of the ISGF3 transcription factor complex. Although early reports indicate Stat3 activation by EGF and IL-6, it has been shown that Stat3 β appears to be activated by both while Stat3 α is activated by EGF, but not by IL-6. Highest expression of Stat4 is seen in testis and myeloid cells. IL-12 has been identified as an activator of Stat4. Stat5 has been shown to be activated by prolactin and by IL-3. Stat6 is involved in IL-4 activated signaling pathways. Synonyms: APRF, Acute-phase response factor, STAT-3, Signal transducer and activator of transcription 3

Molecular Weight:	approx. 88 kDa
Gene ID:	6774
NCBI Accession:	NP_003141
UniProt:	P40763
Pathways:	JAK-STAT Signaling , RTK Signaling , Interferon-gamma Pathway , Neurotrophin Signaling Pathway , Dopaminergic Neurogenesis , Response to Growth Hormone Stimulus , Carbohydrate Homeostasis , Stem Cell Maintenance , Hepatitis C , Protein targeting to Nucleus , Feeding Behaviour , CXCR4-mediated Signaling Events , Signaling of Hepatocyte Growth Factor Receptor

Application Details

Application Notes:	ELISA: 1:10000approx. 1:20000. WB: 1:500approx. 1:1000. IHC: 1:50approx. 1:200. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Restrictions:	For Research Use only

Handling

Concentration:	1,0 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.2., 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	DO NOT FREEZE!

Handling

Storage:	4 °C
Storage Comment:	Store the antibody undiluted at 2-8 °C.

Images




Image 1.

Image 2.