antibodies .- online.com





DDX19B Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image



Go to Product page

| () () () () | 11011 |
|-------------|-------------------------|
| Over | \vee IC \vee \vee |
| 0.0. | |

| Overview | |
|-------------------------------|---|
| Quantity: | 20 μg |
| Target: | DDX19B |
| Protein Characteristics: | Transcript Variant 1 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This DDX19B protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |
| Product Details | |
| Characteristics: | Recombinant human DDX19B (transcript variant 1) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| Target Details | |
| Target: | DDX19B |
| Alternative Name: | Ddx19b (DDX19B Products) |
| Background: | DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative |
| | RNA helicases. They are implicated in a number of cellular processes involving alteration of |
| | RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and |
| | ribosome and spliceosome assembly. Based on their distribution patterns, some members of |

Target Details

this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. This gene encodes a DEAD box protein, which exhibits RNA-dependent ATPase and ATP-dependent RNA-unwinding activities. This protein is recruited to the cytoplasmic fibrils of the nuclear pore complex, where it participates in the export of mRNA from the nucleus. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Molecular Weight:

53.7 kDa

NCBI Accession:

NP_009173

Application Details

Application Notes: Recombinant human proteins can be used for:

Native antigens for optimized antibody production

Positive controls in ELISA and other antibody assays


Comment: The tag is located at the C-terminal.

Restrictions: For Research Use only

Handling

| Concentration: | 50 μg/mL |
|------------------|---|
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze |

immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot