antibodies -online.com





Deoxyguanosine Kinase Protein (DGUOK) (Transcript Variant 1) (Myc-DYKDDDK Tag)



Go to Product pag

1 Image

Overview	
Quantity:	20 μg
Target:	Deoxyguanosine Kinase (DGUOK)
Protein Characteristics:	Transcript Variant 1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Deoxyguanosine Kinase protein is labelled with Myc-DYKDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	Recombinant human Deoxyguanosine kinase (transcript variant 1) protein expressed in
	HEK293 cells. • Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	Deoxyguanosine Kinase (DGUOK)
Alternative Name:	Deoxyguanosine Kinase (DGUOK Products)
Background:	In mammalian cells, the phosphorylation of purine deoxyribonucleosides is mediated
	predominantly by two deoxyribonucleoside kinases, cytosolic deoxycytidine kinase and
	mitochondrial deoxyguanosine kinase. The protein encoded by this gene is responsible for


Target Details

Storage Comment:

Target Details	
	phosphorylation of purine deoxyribonucleosides in the mitochondrial matrix. In addition, this protein phosphorylates several purine deoxyribonucleoside analogs used in the treatment of lymphoproliferative disorders, and this phosphorylation is critical for the effectiveness of the analogs. Alternative splice variants encoding different protein isoforms have been described for this gene.
Molecular Weight:	27.9 kDa
NCBI Accession:	NP_550438
Pathways:	Ribonucleoside Biosynthetic Process
Application Details	
Application Notes:	Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only
Handling	
Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
	O

Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze

immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot