antibodies -online.com





Doublecortin Protein (DCX) (Transcript Variant 4) (Myc-DYKDDDK Tag)



Image



Go to Product pag

\sim				
	$ V \cap$	r\/I	19	٨

Overview		
Quantity:	20 μg	
Target:	Doublecortin (DCX)	
Protein Characteristics:	Transcript Variant 4	
Origin:	Human	
Source:	HEK-293 Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This Doublecortin protein is labelled with Myc-DYKDDDDK Tag.	
Application:	Antibody Production (AbP), Standard (STD)	
Product Details		
Characteristics:	 Recombinant human Doublecortin (transcript variant 4) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone 	
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining	
Target Details		
Target:	Doublecortin (DCX)	
Alternative Name:	Doublecortin (DCX Products)	
Background:	This gene encodes a member of the doublecortin family. The protein encoded by this gene is a	
	cytoplasmic protein and contains two doublecortin domains, which bind microtubules. In the	
	developing cortex, cortical neurons must migrate over long distances to reach the site of their	
	final differentiation. The encoded protein appears to direct neuronal migration by regulating the	

Target Details

organization and stability of microtubules. In addition, the encoded protein interacts with LIS1, the regulatory gamma subunit of platelet activating factor acetylhydrolase, and this interaction is important to proper microtubule function in the developing cortex. Mutations in this gene cause abnormal migration of neurons during development and disrupt the layering of the cortex, leading to epilepsy, mental retardation, subcortical band heterotopia ("double cortex" syndrome) in females and lissencephaly ("smooth brain" syndrome) in males. Multiple transcript variants encoding different isoforms have been found for this gene.

Molecular Weight:

39.9 kDa

NCBI Accession:

NP_835364

Application Details

Application Notes:

Recombinant human proteins can be used for:

Native antigens for optimized antibody production

Positive controls in ELISA and other antibody assays

Comment:

The tag is located at the C-terminal.

Restrictions:

For Research Use only

Handling

50 μg/mL

Buffer:

25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.


Storage:

-80 °C

Storage Comment:

Store at -80 $^{\circ}\text{C}.$ Thaw on ice, aliquot to individual single-use tubes, and then re-freeze

immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot