

Datasheet for ABIN2719686

DCLK2 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)[Go to Product page](#)

Overview

| | |
|-------------------------------|-------------------------------------------------------|
| Quantity: | 20 µg |
| Target: | DCLK2 |
| Protein Characteristics: | Transcript Variant 1 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This DCLK2 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |

Product Details

| | |
|------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Characteristics: | <ul style="list-style-type: none">• Recombinant human Doublecortin-like kinase 2 (transcript variant 1) protein expressed in HEK293 cells.• Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |

Target Details

| | |
|-------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Target: | DCLK2 |
| Alternative Name: | Doublecortin-Like Kinase 2 (DCLK2 Products) |
| Background: | <p>This gene encodes a member of the protein kinase superfamily and the doublecortin family.</p> <p>The protein encoded by this gene contains two N-terminal doublecortin domains, which bind microtubules and regulate microtubule polymerization, a C-terminal serine/threonine protein</p> |

Target Details

kinase domain, which shows substantial homology to Ca²⁺/calmodulin-dependent protein kinase, and a serine/proline-rich domain in between the doublecortin and the protein kinase domains, which mediates multiple protein-protein interactions. The microtubule-polymerizing activity of the encoded protein is independent of its protein kinase activity. Mouse studies show that the DCX gene, another family member, and this gene share function in the establishment of hippocampal organization and that their absence results in a severe epileptic phenotype and lethality, as described in human patients with lissencephaly. Multiple alternatively spliced transcript variants have been identified.

| | |
|-------------------|------------------------------|
| Molecular Weight: | 83.4 kDa |
| NCBI Accession: | NP_001035350 |

Application Details

| | |
|--------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|
| Application Notes: | Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |

Handling

| | |
|------------------|---------------------------------------------------------------------------------------------------------------------------------------------------|
| Concentration: | 50 µg/mL |
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |