# antibodies -online.com





## **DPYS Protein (Myc-DYKDDDDK Tag)**



Image



Go to Product page

| $\sim$ |     |      |            |
|--------|-----|------|------------|
|        | N/6 | 1//r | $I \cap V$ |

| Quantity:                     | 20 μg   |
|-------------------------------|---|
| Target:                       | DPYS  |
| Origin:                       | Human   |
| Source:                       | HEK-293 Cells   |
| Protein Type:                 | Recombinant   |
| Purification tag / Conjugate: | This DPYS protein is labelled with Myc-DYKDDDDK Tag.  |
| Application:                  | Antibody Production (AbP), Standard (STD)   |
| Product Details               |   |
| Characteristics:              | <ul> <li>Recombinant human DPYS protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>  |
| Purity:                       | > 80 % as determined by SDS-PAGE and Coomassie blue staining  |
| Target Details                |   |
| Target:                       | DPYS  |
| Alternative Name:             | Dpys (DPYS Products)  |
| Background:                   | Dihydropyrimidinase catalyzes the conversion of 5,6-dihydrouracil to 3-ureidopropionate in pyrimidine metabolism. Dihydropyrimidinase is expressed at a high level in liver and kidney as a major 2.5-kb transcript and a minor 3.8-kb transcript. Defects in the DPYS gene are linked to dihydropyrimidinuria. |
| Molecular Weight:             | 56.4 kDa  |

#### **Target Details**

| NCBI Accession: NP_001376 | NCBI Accession: | NP_ | _001376 |
|---------------------------|-----------------|-----|---------|
|---------------------------|-----------------|-----|---------|


### **Application Details**

| Application Notes: | Recombinant human proteins can be used for:          |
|--------------------|--|
|                    | Native antigens for optimized antibody production    |
|                    | Positive controls in ELISA and other antibody assays |
| Comment:           | The tag is located at the C-terminal.                |
| Restrictions:      | For Research Use only                                |

#### Handling

| Concentration:   | 50 μg/mL  |
|------------------|---|
| Buffer:          | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.  |
| Storage:         | -80 °C  |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

#### Images



#### **Western Blotting**

Image 1. Validation with Western Blot