antibodies .- online.com





DUSP13 Protein (Transcript Variant 6) (Myc-DYKDDDDK Tag)



Image



-						
O	V	e	rv	1	е	W

20 μg	
DUSP13	
Transcript Variant 6	
Human	
HEK-293 Cells	
Recombinant	
This DUSP13 protein is labelled with Myc-DYKDDDDK Tag.	
Antibody Production (AbP), Standard (STD)	
 Recombinant human DUSP13 (transcript variant 6) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone 	
> 80 % as determined by SDS-PAGE and Coomassie blue staining	
DUSP13	
usp13 (DUSP13 Products)	
Members of the protein-tyrosine phosphatase superfamily cooperate with protein kinases to regulate cell proliferation and differentiation. This superfamily is separated into two families based on the substrate that is dephosphorylated. One family, the dual specificity phosphatases (DSPs) acts on both phosphotyrosine and phosphoserine/threonine residues. This gene	

encodes different but related DSP proteins through the use of non-overlapping open reading frames, alternate splicing, and presumed different transcription promoters. Expression of the distinct proteins from this gene has been found to be tissue specific and the proteins may be involved in postnatal development of specific tissues. A protein encoded by the upstream ORF was found in skeletal muscle, whereas the encoded protein from the downstream ORF was found only in testis. In mouse, a similar pattern of expression was found. Multiple alternatively spliced transcript variants were described, but the full-length sequence of only some were determined.

Molecular Weight:

22 kDa

NCBI Accession:

NP 057448

Application Details

Application Notes:

Recombinant human proteins can be used for:

Native antigens for optimized antibody production

Positive controls in ELISA and other antibody assays

Comment:

The tag is located at the C-terminal.

Restrictions:


For Research Use only

Handling

Concentration:	50 μg/mL			
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.			
Storage:	-80 °C			

Storage Comment:

Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot