
antibodies -online.com





DTNBP1 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image



Go to Product page

| Overview | |
|-------------------------------|--|
| Quantity: | 20 μg |
| Target: | DTNBP1 |
| Protein Characteristics: | Transcript Variant 1 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This DTNBP1 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |
| Product Details | |
| Characteristics: | Recombinant human Dysbindin (transcript variant 1) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| Target Details | |
| Target: | DTNBP1 |
| Alternative Name: | Dysbindin (DTNBP1 Products) |
| Background: | This gene encodes a protein that may play a role in organelle biogenesis associated with |

melanosomes, platelet dense granules, and lysosomes. A similar protein in mouse is a

component of a protein complex termed biogenesis of lysosome-related organelles complex 1 (BLOC-1), and binds to alpha- and beta-dystrobrevins, which are components of the dystrophin-

Target Details

| | associated protein complex (DPC). Mutations in this gene are associated with Hermansky-Pudlak syndrome type 7. This gene may also be associated with schizophrenia. Multiple transcript variants encoding distinct isoforms have been identified for this gene. |
|-------------------|---|
| Molecular Weight: | 39.3 kDa |
| NCBI Accession: | NP_115498 |
| Pathways: | Synaptic Membrane, Regulation of G-Protein Coupled Receptor Protein Signaling |


Application Details

| Application Notes: | Recombinant human proteins can be used for: |
|--------------------|--|
| | Native antigens for optimized antibody production |
| | Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |

Handling

| Concentration: | 50 μg/mL |
|------------------|---|
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |

Images



Western Blotting

Image 1. Validation with Western Blot