

Datasheet for ABIN272005
anti-CD130/gp130 antibody



[Go to Product page](#)

2 Images

Overview

Quantity:	0.1 mg
Target:	CD130/gp130 (IL6ST)
Reactivity:	Human, Mouse, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CD130/gp130 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Specificity:	This antibody detects endogenous levels of CD130/gp130 protein. (region surrounding Ser778)
Cross-Reactivity (Details):	Species reactivity (tested): Human, Mouse.
Purification:	Affinity Chromatography using epitope-specific immunogen.

Target Details

Target:	CD130/gp130 (IL6ST)
Alternative Name:	CD130 / IL6ST (IL6ST Products)
Background:	IL-6 activates intracellular signaling through binding a receptor consisting of an 80 kDa ligand-binding protein (IL-6R) and a second protein of 130 kDa. IL-6 first binds to IL-6R which subsequently associates with a gp130 dimer. The active signaling complex consists of at minimum IL-6, IL-6R and a dimer of two gp130 proteins that are linked by a disulfide bond. A

Target Details

soluble form of IL-6R is generated by proteolytic cleavage of the membrane-bound precursor and can function as an agonistic molecule that can actively participate in cell-to-cell signaling. The second subunit of the IL-6 complex, gp130, also functions as a component of several additional receptor complexes including leukemia inhibitory factor (LIF), oncostatin M (OSM), ciliary neurotrophic factor (CNTF) and IL-11. LIF binds to the LIF receptor with low affinity and to a complex of the LIF receptor and gp130 with high affinity, while OSM appears to bind to gp130 with low affinity and to a complex of gp130 and the LIF receptor with high affinity. Synonyms: IL-6R-beta, IL6 receptor beta, Interleukin-6 receptor subunit beta

Molecular Weight:	approx. 160 kDa
Gene ID:	3572
NCBI Accession:	NP_002175
UniProt:	P40189
Pathways:	JAK-STAT Signaling , Cellular Glucan Metabolic Process , Autophagy , Smooth Muscle Cell Migration , Cancer Immune Checkpoints

Application Details

Application Notes:	ELISA: 1: 20000approx. 1: 40000. IHC: 1: 50approx. 1: 200. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Restrictions:	For Research Use only

Handling

Concentration:	1.0 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH ~7.2, 0.05 % Sodium Azide
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freezing and thawing.
Storage:	4 °C/-20 °C
Storage Comment:	Store the antibody undiluted at 2-8 °C for one month or (in aliquots) at -20 °C for longer.




Image 1.




Image 2.