
antibodies .- online.com







EPH Receptor A5 Protein (EPHA5) (DYKDDDDK Tag)



Image



| <i>ا</i> ۱ | 1 | - | M | /1 | 0 | |
|------------|-----|----------|-----|-----|--------|------|
| u | 1// | \vdash | 1 \ | / I | - | \/\/ |
| \sim | v | \sim | r٧ | | \sim | v v |

| Overview | | | |
|-------------------------------|-----------------------------------------------------------------------------------------------|--|--|
| Quantity: | 20 μg | | |
| Target: | EPH Receptor A5 (EPHA5) | | |
| Origin: | Human | | |
| Source: | Insect cells (Sf9) | | |
| Protein Type: | Recombinant | | |
| Purification tag / Conjugate: | This EPH Receptor A5 protein is labelled with DYKDDDDK Tag. | | |
| Application: | Antibody Production (AbP), Standard (STD) | | |
| Product Details | | | |
| Characteristics: | Recombinant human EPHA5 (C-term DDK tag, transcript variant 1) protein expressed in Sf9 | | |
| | cells. • Produced with end-sequenced ORF clone | | |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining | | |
| Target Details | | | |
| Target: | EPH Receptor A5 (EPHA5) | | |
| Alternative Name: | Epha5 (EPHA5 Products) | | |
| Background: | This gene belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family. EPH | | |
| | and EPH-related receptors have been implicated in mediating developmental events, | | |
| | particularly in the nervous system. Receptors in the EPH subfamily typically have a single | | |
| | kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type | | |

Storage:


Storage Comment:

| Target Details | | | |
|---------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--|
| | III repeats. The ephrin receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Alternatively spliced transcript variants encoding different isoforms have been described. | | |
| Molecular Weight: | 60.8 kDa | | |
| NCBI Accession: | NP_004430 | | |
| Pathways: | RTK Signaling | | |
| Application Details | | | |
| Application Notes: | Recombinant human proteins can be used for: Native antigens for optimized antibody production Positive controls in ELISA and other antibody assays | | |
| Comment: | The tag is located at the C-terminal. | | |
| Restrictions: | For Research Use only | | |
| Handling | | | |
| Concentration: | 50 μg/mL | | |
| Buffer: | 50 mM Tris-HCl, pH 8.0, 100 mM glycine, 10 % glycerol. Store at -80C. Avoid repeated freeze-thaw cycles. Stable for at least 3 months from receipt of products under proper storage and handling conditions. | | |

-80 °C

Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze

immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot