# antibodies -online.com





# **ERBB4 Protein (Myc-DYKDDDDK Tag)**

2 Images




Go to Product page

$\sim$				
/ \	1/0	r\ /	01	Α.
	$\vee$	r 🗤	$\square$	Λ

O V CI VICVV		
Quantity:	20 μg	
Target:	ERBB4	
Origin:	Human	
Source:	HEK-293 Cells	
Protein Type:	Recombinant	
Biological Activity:	Active	
Purification tag / Conjugate:	This ERBB4 protein is labelled with Myc-DYKDDDDK Tag.	
Application:	Antibody Production (AbP), Standard (STD), Functional Studies (Func), Protein Interaction (PI)	
Product Details		
Specificity:	Optimal preservation of protein structure, post-translational modifications and functions.	
Characteristics:	<ul> <li>Recombinant human ERBB4 / HER4 (transcript variant JM-a/CVT-2) protein expressed in HEK293 cells.</li> </ul>	
	Produced with end-sequenced ORF clone	
	Tested for bioactivity.	
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining	
Biological Activity Comment:	ErbB4 activity verified in a biochemical assay:,ErbB4 activity verified in a biochemical assay:	
Target Details		
Target:	ERBB4	
Alternative Name:	Erbb4,her4 (ERBB4 Products)	


## **Target Details**

•			
Background:	This gene is a member of the Tyr protein kinase family and the epidermal growth factor		
	receptor subfamily. It encodes a single-pass type I membrane protein with multiple cysteine		
	rich domains, a transmembrane domain, a tyrosine kinase domain, a phosphotidylinositol-3		
	kinase binding site and a PDZ domain binding motif. The protein binds to and is activated by		
	neuregulins and other factors and induces a variety of cellular responses including mitogenesis		
	and differentiation. Multiple proteolytic events allow for the release of a cytoplasmic fragment		
	and an extracellular fragment. Mutations in this gene have been associated with cancer.		
	Alternatively spliced variants which encode different protein isoforms have been described		
	however, not all variants have been fully characterized.		
Molecular Weight:	142.6 kDa		
NCBI Accession:	NP_001036064		
Pathways:	RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin		
	Signaling Pathway		
Application Details			
Application Notes:	Recombinant human proteins can be used for:		
	Native antigens for optimized antibody production		
	Positive controls in ELISA and other antibody assays		
	Protein-protein interaction		
	In vitro biochemical assays and cell-based functional assays		
Comment:	The tag is located at the C-terminal.		
Restrictions:	For Research Use only		
Handling			
Concentration:	> 50 μg/mL		
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.		
Storage:	-80 °C		
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze		
	immediately. Only 2-3 freeze thaw cycles are recommended.		



### **Activity Assay**

Image 1. Bioactivity measured with Activity Assay



#### **Western Blotting**

Image 2. Validation with Western Blot