## antibodies.com

## Datasheet for ABIN2721250 FMO3 Protein (Transcript Variant 2) (Myc-DYKDDDDK Tag)



Image

## Overview


Quantity:	20 µg
Target:	FM03
Protein Characteristics:	Transcript Variant 2
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This FMO3 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	<ul> <li>Recombinant human FMO3 (transcript variant 2) protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	FM03
Alternative Name:	Fmo3 (FMO3 Products)
Background:	Flavin-containing monooxygenases (FMO) are an important class of drug-metabolizing
	enzymes that catalyze the NADPH-dependent oxygenation of various nitrogen-,sulfur-, and
	phosphorous-containing xenobiotics such as therapeutic drugs, dietary compounds, pesticides,

and other foreign compounds. The human FMO gene family is composed of 5 genes and

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN2721250 | 09/11/2023 | Copyright antibodies-online. All rights reserved.

	multiple pseudogenes. FMO members have distinct developmental- and tissue-specific
	expression patterns. The expression of this FMO3 gene, the major FMO expressed in adult liver,
	can vary up to 20-fold between individuals. This inter-individual variation in FMO3 expression
	levels is likely to have significant effects on the rate at which xenobiotics are metabolised and,
	therefore, is of considerable interest to the pharmaceutical industry. This transmembrane
	protein localizes to the endoplasmic reticulum of many tissues. Alternative splicing of this gene
	results in multiple transcript variants encoding different isoforms. Mutations in this gene cause
	the disorder trimethylaminuria (TMAu) which is characterized by the accumulation and
	excretion of unmetabolized trimethylamine and a distinctive body odor. In healthy individuals,
	trimethylamine is primarily converted to the non odorous trimethylamine N-oxide.[provided by
	RefSeq, Jan 2016].
Molecular Weight:	59.9 kDa
NCBI Accession:	NP_001002294
Application Details	
Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only
Handling	
Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze
	immediately. Only 2-3 freeze thaw cycles are recommended.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN2721250 | 09/11/2023 | Copyright antibodies-online. All rights reserved.



Western Blotting

Image 1. Validation with Western Blot

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN2721250 | 09/11/2023 | Copyright antibodies-online. All rights reserved.