antibodies .- online.com





GIT2 Protein (Transcript Variant 4) (Myc-DYKDDDDK Tag)



Image



)\	'e	r٧	/ (91	Ν

Overview	
Quantity:	20 μg
Target:	GIT2
Protein Characteristics:	Transcript Variant 4
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This GIT2 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	 Recombinant human G protein-coupled receptor kinase interacting ArfGAP 2 (GIT2), transcript variant 4 (transcript variant 4) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	GIT2
Alternative Name:	G Protein-Coupled Receptor Kinase Interacting Arfgap 2 (Git2) (GIT2 Products)
Background:	This gene encodes a member of the GIT protein family, which interact with G protein-coupled receptor kinases and possess ADP-ribosylation factor (ARF) GTPase-activating protein (GAP) activity. GIT proteins traffic between cytoplasmic complexes, focal adhesions, and the cell

periphery, and interact with Pak interacting exchange factor beta (PIX) to form large oligomeric
complexes that transiently recruit other proteins. GIT proteins regulate cytoskeletal dynamics
and participate in receptor internalization and membrane trafficking. This gene has been shown
to repress lamellipodial extension and focal adhesion turnover, and is thought to regulate cell
motility. This gene undergoes extensive alternative splicing to generate multiple isoforms, but
the full-length nature of some of these variants has not been determined. The various isoforms
have functional differences, with respect to ARF GAP activity and to G protein-coupled receptor
kinase 2 binding.


Molecular Weight:	52.4 kDa
NCBI Accession:	NP_631940
Pathways: Regulation of G-Protein Coupled Receptor Protein Signaling	

Application Details

Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only

Handling

Concentration:	50 μg/mL
Buffer: 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.	
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot