
## antibodies -online.com





## GCK Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image




Go to Product page

Overview	
Quantity:	20 μg
Target:	GCK
Protein Characteristics:	Transcript Variant 1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This GCK protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	<ul> <li>Recombinant human Glucokinase / Hexokinase-4 (transcript variant 1) protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	GCK
Alternative Name:	Glucokinase,hexokinase-4 (GCK Products)
Background:	Hexokinases phosphorylate glucose to produce glucose-6-phosphate, the first step in most glucose metabolism pathways. Alternative splicing of this gene results in three tissue-specific

forms of glucokinase, one found in pancreatic islet beta cells and two found in liver. The protein

## **Target Details**

Target Details	
	localizes to the outer membrane of mitochondria. In contrast to other forms of hexokinase, this enzyme is not inhibited by its product glucose-6-phosphate but remains active while glucose is abundant. Mutations in this gene have been associated with non-insulin dependent diabetes mellitus (NIDDM), maturity onset diabetes of the young, type 2 (MODY2) and persistent hyperinsulinemic hypoglycemia of infancy (PHHI).
Molecular Weight:	52 kDa
NCBI Accession:	NP_000153
Pathways:	MAPK Signaling, Positive Regulation of Peptide Hormone Secretion, Carbohydrate Homeostasis, Cellular Glucan Metabolic Process, Regulation of Carbohydrate Metabolic Process
Application Details	
Application Notes:	Recombinant human proteins can be used for:  Native antigens for optimized antibody production  Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only
Handling	
Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.



## **Western Blotting**

Image 1. Validation with Western Blot