antibodies - online.com





GSTM2 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Overview

Image



Go to Product page

Quantity:	20 μg
Target:	GSTM2
Protein Characteristics:	Transcript Variant 1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This GSTM2 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	

• Recombinant human GSTM2 / GST4 (transcript variant 1) protein expressed in HEK293 cells.

Product Details

Characteristics:

Purity:

Target Details	
Target:	GSTM2
Alternative Name:	Gstm2,gst4 (GSTM2 Products)
Background:	Cytosolic and membrane-bound forms of glutathione S-transferase are encoded by two distinct supergene families. At present, eight distinct classes of the soluble cytoplasmic mammalian glutathione S-transferases have been identified: alpha, kappa, mu, omega, pi, sigma, theta and zeta. This gene encodes a glutathione S-transferase that belongs to the mu class. The mu class

> 80 % as determined by SDS-PAGE and Coomassie blue staining

· Produced with end-sequenced ORF clone

Storage:


Storage Comment:

-80 °C

Target Details	
	of enzymes functions in the detoxification of electrophilic compounds, including carcinogens, therapeutic drugs, environmental toxins and products of oxidative stress, by conjugation with glutathione. The genes encoding the mu class of enzymes are organized in a gene cluster on chromosome 1p13.3 and are known to be highly polymorphic. These genetic variations can change an individual's susceptibility to carcinogens and toxins as well as affect the toxicity and efficacy of certain drugs.
Molecular Weight:	25.6 kDa
NCBI Accession:	NP_000839
Pathways:	Negative Regulation of Transporter Activity
Application Details	
Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only
Handling	
Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.

immediately. Only 2-3 freeze thaw cycles are recommended.

Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze



Western Blotting

Image 1. Validation with Western Blot