antibodies - online.com







HAAO Protein (Myc-DYKDDDDK Tag)



Image



\sim				
	$ V \cap$	r\/I	19	٨

20 μg
HAAO
Human
HEK-293 Cells
Recombinant
This HAAO protein is labelled with Myc-DYKDDDDK Tag.
Antibody Production (AbP), Standard (STD)
 Recombinant human HAAO protein expressed in HEK293 cells. Produced with end-sequenced ORF clone
> 80 % as determined by SDS-PAGE and Coomassie blue staining
HAAO
Haao (HAAO Products)
3-Hydroxyanthranilate 3,4-dioxygenase is a monomeric cytosolic protein belonging to the family of intramolecular dioxygenases containing nonheme ferrous iron. It is widely distributed in peripheral organs, such as liver and kidney, and is also present in low amounts in the central nervous system. HAAO catalyzes the synthesis of quinolinic acid (QUIN) from 3-hydroxyanthranilic acid. QUIN is an excitotoxin whose toxicity is mediated by its ability to

Target Details

	activate glutamate N-methyl-D-aspartate receptors. Increased cerebral levels of QUIN may
	participate in the pathogenesis of neurologic and inflammatory disorders. HAAO has been
	suggested to play a role in disorders associated with altered tissue levels of QUIN.
Molecular Weight:	32.4 kDa
NCBI Accession:	NP_036337


Application Details

Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only

Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.

Images



Western Blotting

Image 1. Validation with Western Blot