antibodies -online.com





HAX1 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image




Go to Product page

| Overview | |
|-------------------------------|--|
| Quantity: | 20 μg |
| Target: | HAX1 |
| Protein Characteristics: | Transcript Variant 1 |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This HAX1 protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |
| Product Details | |
| Characteristics: | Recombinant human HAX1 (transcript variant 1) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| Target Details | |
| Target: | HAX1 |
| Alternative Name: | Hax1 (HAX1 Products) |
| Background: | The protein encoded by this gene is known to associate with hematopoietic cell-specific Lyn substrate 1, a substrate of Src family tyrosine kinases. It also interacts with the product of the |

polycystic kidney disease 2 gene, mutations in which are associated with autosomal-dominant polycystic kidney disease, and with the F-actin-binding protein, cortactin. It was earlier thought

Target Details

| | that this gene product is mainly localized in the mitochondria, however, recent studies indicate it to be localized in the cell body. Mutations in this gene result in autosomal recessive severe congenital neutropenia, also known as Kostmann disease. Two transcript variants encoding different isoforms have been found for this gene. |
|---------------------|--|
| Molecular Weight: | 31.4 kDa |
| NCBI Accession: | NP_006109 |
| Pathways: | Regulation of Actin Filament Polymerization |
| Application Details | |
| Application Notes: | Recombinant human proteins can be used for: |
| | Native antigens for optimized antibody production |
| | Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |
| Handling | |
| Concentration: | 50 μg/mL |
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. |
| Storage: | -80 °C |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended. |



Western Blotting

Image 1. Validation with Western Blot