

Datasheet for ABIN2722630

HGF Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)[Go to Product page](#)[1 Image](#)[1 Publication](#)

Overview

Quantity:	20 µg
Target:	HGF
Protein Characteristics:	Transcript Variant 1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This HGF protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)

Product Details

Characteristics:	<ul style="list-style-type: none">• Recombinant human Hepatocyte growth factor / HGF (transcript variant 1) protein expressed in HEK293 cells.• Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining

Target Details

Target:	HGF
Alternative Name:	Hepatocyte Growth Factor,hgf (HGF Products)
Background:	This gene encodes a protein that binds to the hepatocyte growth factor receptor to regulate cell growth, cell motility and morphogenesis in numerous cell and tissue types. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is

Target Details

proteolytically processed to generate alpha and beta chains, which form the mature heterodimer. This protein is secreted by mesenchymal cells and acts as a multi-functional cytokine on cells of mainly epithelial origin. This protein also plays a role in angiogenesis, tumorigenesis, and tissue regeneration. Although the encoded protein is a member of the peptidase S1 family of serine proteases, it lacks peptidase activity. Mutations in this gene are associated with nonsyndromic hearing loss.

Molecular Weight: 79.6 kDa

NCBI Accession: [NP_000592](#)

Pathways: [RTK Signaling](#), [Carbohydrate Homeostasis](#), [Glycosaminoglycan Metabolic Process](#), [Synaptic Membrane](#), [Signaling of Hepatocyte Growth Factor Receptor](#)

Application Details

Application Notes: Recombinant human proteins can be used for:
Native antigens for optimized antibody production
Positive controls in ELISA and other antibody assays

Comment: The tag is located at the C-terminal.

Restrictions: For Research Use only

Handling

Concentration: 50 µg/mL


Buffer: 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.

Storage: -80 °C

Storage Comment: Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.

Publications

Product cited in: Ueda, Ito, Shiraishi, Taniguchi, Kayukawa, Nakanishi, Nakamura, Naya, Hongo, Kamoi, Okihara, Kawauchi, Miki: "PAX2 promoted prostate cancer cell invasion through transcriptional regulation of HGF in an in vitro model." in: **Biochimica et biophysica acta**, Vol. 1852, Issue 11, pp. 2467-73, (2015) ([PubMed](#)).



Western Blotting

Image 1. Validation with Western Blot