antibodies -online.com





Histone H2B Type 1-C/E/F/G/I (LOC100407767) protein (Myc-DYKDDDK Tag)



Go to Product pac

1 Image

Overview	
Quantity:	20 μg
Target:	Histone H2B Type 1-C/E/F/G/I (LOC100407767)
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	Myc-DYKDDDDK Tag
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	 Recombinant human Histone H2B type 1-C/E/F/G/I protein expressed in HEK293 cells. Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	Histone H2B Type 1-C/E/F/G/I (LOC100407767)
Alternative Name:	Histone h2b Type 1-C/e/f/g/i (LOC100407767 Products)
Background:	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the
	chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B,
	H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in
	repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between
	nucleosomes and functions in the compaction of chromatin into higher order structures. This

Target Details

gene is intronless and encodes a replication-dependent histone that is a member of the histone
H2B family. The protein has antibacterial and antifungal antimicrobial activity. Transcripts from
this gene lack polyA tails but instead contain a palindromic termination element. This gene is
found in the large histone gene cluster on chromosome 6.

Molecular Weight:	13.7 kDa

NCBI Accession: NP_003513


Application Details

Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only

Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.

Images



Western Blotting

Image 1. Validation with Western Blot