# antibodies -online.com







# HIST1H2BM Protein (Myc-DYKDDDDK Tag)



Image



$C_0 + c$	Produc	+
	Promi	บ กลกย

( )	ve	K\ /		A .
	$\cup$	1 V/	Щ.	V۷

Overview	
Quantity:	20 μg
Target:	HIST1H2BM
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This HIST1H2BM protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	<ul> <li>Recombinant human Histone H2B type 1-M protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	HIST1H2BM
Abstract:	HIST1H2BM Products
Background:	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the
	chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B,
	H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in
	repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between
	nucleosomes and functions in the compaction of chromatin into higher order structures. This

## **Target Details**

gene is intronless and encodes a replication-dependent histone that is a member of the histone
H2B family. Transcripts from this gene lack polyA tails but instead contain a palindromic
termination element. This gene is found in the small histone gene cluster on chromosome
6p22-p21.3.
10.010.

Molecular Weight: 13.8 kDa

NCBI Accession: NP\_003512


# **Application Details**

Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only

# Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.

#### **Images**



## **Western Blotting**

Image 1. Validation with Western Blot