antibodies .- online.com





HRAS Protein (Transcript Variant 3) (Myc-DYKDDDDK Tag)



Image

Characteristics:



Overview	
Quantity:	20 μg
Target:	HRAS
Protein Characteristics:	Transcript Variant 3
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This HRAS protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	

Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	HRAS
Alternative Name:	Hras (HRAS Products)
Background:	This gene belongs to the Ras oncogene family, whose members are related to the transforming

· Produced with end-sequenced ORF clone

genes of mammalian sarcoma retroviruses. The products encoded by these genes function in signal transduction pathways. These proteins can bind GTP and GDP, and they have intrinsic GTPase activity. This protein undergoes a continuous cycle of de- and re-palmitoylation, which

• Recombinant human HRAS (transcript variant 3) protein expressed in HEK293 cells.

Multiple transcript variants, which encode different isoforms, have been identified for this gene.
cancers, including bladder cancer, follicular thyroid cancer, and oral squamous cell carcinoma.
appearance and cardiovascular abnormalities. Defects in this gene are implicated in a variety of
formation, mental retardation, skin and musculoskeletal abnormalities, distinctive facial
at the prenatal stage, growth deficiency at the postnatal stage, predisposition to tumor
Mutations in this gene cause Costello syndrome, a disease characterized by increased growth
regulates its rapid exchange between the plasma membrane and the Golgi apparatus.

Molecular Weight: 21.1 kDa

NCBI Accession: NP_001123914

Pathways:


p53 Signaling, MAPK Signaling, RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR
Signaling Pathway, Neurotrophin Signaling Pathway, Hepatitis C, Autophagy, Signaling Events
mediated by VEGFR1 and VEGFR2, Signaling of Hepatocyte Growth Factor Receptor,
Regulation of long-term Neuronal Synaptic Plasticity, VEGF Signaling, BCR Signaling

Application Details

Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only

Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot