

#### Datasheet for ABIN2723367

# IL17RC Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image



Go to Product page

Overview	

Quantity:	20 μg
Target:	IL17RC
Protein Characteristics:	Transcript Variant 1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL17RC protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	<ul> <li>Recombinant human IL17RC (transcript variant 1) protein expressed in HEK293 cells.</li> <li>Produced with end-sequenced ORF clone</li> </ul>
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	IL17RC
Alternative Name:	II17rc (IL17RC Products)
Background:	This gene encodes a single-pass type I membrane protein that shares similarity with the interleukin-17 receptor (IL-17RA). Unlike IL-17RA, which is predominantly expressed in hemopoietic cells, and binds with high affinity to only IL-17A, this protein is expressed in nonhemopoietic tissues, and binds both IL-17A and IL-17F with similar affinities. The

# **Target Details**

proinflammatory cytokines, IL-17A and IL-17F, have been implicated in the progression of
inflammatory and autoimmune diseases. Multiple alternatively spliced transcript variants
encoding different isoforms have been detected for this gene, and it has been proposed that
soluble, secreted proteins lacking transmembrane and intracellular domains may function as
extracellular antagonists to cytokine signaling.

Molecular Weight:

83.9 kDa

NCBI Accession:

NP\_703191

# **Application Details**


Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.

Restrictions:

For Research Use only

# Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.



# **Western Blotting**

Image 1. Validation with Western Blot