antibodies -online.com





ITGA8 Protein (Myc-DYKDDDDK Tag)





Go to Product page

\sim					
	1//6	r	V I	Θ	Λ

20 μg	
. 5	
ITGA8	
Human	
HEK-293 Cells	
Recombinant	
This ITGA8 protein is labelled with Myc-DYKDDDDK Tag.	
Antibody Production (AbP), Standard (STD)	
 Recombinant human Integrin alpha-8 / ITGA8 protein expressed in HEK293 cells. Produced with end-sequenced ORF clone 	
> 80 % as determined by SDS-PAGE and Coomassie blue staining	
ITGA8	
Integrin alpha-8,itga8 (ITGA8 Products)	
Integrins are heterodimeric transmembrane receptor proteins that mediate numerous cellular processes including cell adhesion, cytoskeletal rearrangement, and activation of cell signaling pathways. Integrins are composed of alpha and beta subunits. This gene encodes the alpha 8 subunit of the heterodimeric integrin alpha8beta1 protein. The encoded protein is a single-pass type 1 membrane protein that contains multiple FG-GAP repeats. This repeat is predicted to	

Target Details

fold into a beta propeller structure. This gene regulates the recruitment of mesenchymal cells into epithelial structures, mediates cell-cell interactions, and regulates neurite outgrowth of sensory and motor neurons. The integrin alpha8beta1 protein thus plays an important role in wound-healing and organogenesis. Mutations in this gene have been associated with renal hypodysplasia/aplasia-1 (RHDA1) and with several animal models of chronic kidney disease. Alternate splicing results in multiple transcript variants encoding distinct isoforms.

Molecular Weight:

117.3 kDa

NCBI Accession:

NP_003629

Application Details

Application Notes: Recombinant human proteins can be used for:

Native antigens for optimized antibody production

Positive controls in ELISA and other antibody assays


Comment: The tag is located at the C-terminal.

Restrictions: For Research Use only

Handling

Concentration:	50 μg/mL	
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.	
Storage: -80 °C		
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze	

immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot