# antibodies - online.com







# **IL-19 Protein (Transcript Variant 2)**



## Image




Goto	Drad	LIGH	2000
(7() (()	PI(00)	UICH	Datte

U	V	er	V	ıе	W

Quantity:	10 μg
Target:	IL-19 (IL19)
Protein Characteristics:	Transcript Variant 2
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Biological Activity:	Active
Application:	Functional Studies (Func), Antibody Production (AbP), Standard (STD), Protein Interaction (PI)
Product Details	
Specificity:	Optimal preservation of protein structure, post-translational modifications and functions.
Characteristics:	<ul> <li>Recombinant human Interleukin-19 / IL19 (transcript variant 2) protein expressed in E. coli.</li> <li>Produced with end-sequenced ORF clone</li> <li>Tested for bioactivity.</li> </ul>
Purity:	> 95 % as determined by SDS-PAGE and Coomassie blue staining
Endotoxin Level:	Endotoxin level is <0.1 ng/μg of protein (<1EU/μg).
Biological Activity Comment:	Determined by its ability to activate STAT following receptor-ligand interaction.
Target Details	
Target:	IL-19 (IL19)

### **Target Details**

Altornative News	Interlaukin 10 il10 (II 10 Products)
Alternative Name:	Interleukin-19,il19 (IL19 Products)
Background:	The protein encoded by this gene is a cytokine that belongs to the IL10 cytokine subfamily. This
	cytokine is found to be preferentially expressed in monocytes. It can bind the IL20 receptor
	complex and lead to the activation of the signal transducer and activator of transcription 3
	(STAT3). A similar cytokine in mouse is reported to up-regulate the expression of IL6 and TNF-
	alpha and induce apoptosis, which suggests a role of this cytokine in inflammatory responses.
	Alternatively spliced transcript variants encoding the distinct isoforms have been described.
Molecular Weight:	17.9 kDa
NCBI Accession:	NP_037503
Application Details	
Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
	Protein-protein interaction
	In vitro biochemical assays and cell-based functional assays
Restrictions:	For Research Use only
Handling	
Buffer:	Lyophilized from a 0.2 µM filtered solution of 20 mM phosphate buffer,100 mM NaCl, pH 7.2
Handling Advice:	Resuspend the protein in the desired concentration in proper buffer
Storage:	-80 °C
Storage Comment:	Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze
	immediately. Only 2-3 freeze thaw cycles are recommended.



### **Western Blotting**

Image 1. Validation with Western Blot