antibodies -online.com





KPNA1 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image



Go to Product page

Overview	V
OVEIVIEV	۷

Overview	
Quantity:	20 μg
Target:	KPNA1
Protein Characteristics:	Transcript Variant 1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This KPNA1 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	Recombinant human KPNA1 / Importin alpha-1 (transcript variant 1) protein expressed in HEK293 cells.
	Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	KPNA1
Alternative Name:	Kpna1,importin alpha-1 (KPNA1 Products)
Background:	Functions in nuclear protein import as an adapter protein for nuclear receptor KPNB1. Binds
	specifically and directly to substrates containing either a simple or bipartite NLS motif. Docking
	of the importin/substrate complex to the nuclear pore complex (NPC) is mediated by KPNB1

through binding to nucleoporin FxFG repeats and the complex is subsequently translocated through the pore by an energy requiring, Ran-dependent mechanism. At the nucleoplasmic side of the NPC, Ran binds to importin-beta and the three components separate and importin-alpha and -beta are re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran from importin. The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus. In vitro, mediates the nuclear import of human cytomegalovirus UL84 by recognizing a non-classical NLS. [UniProtKB/Swiss-Prot Function]


Molecular Weight:	60 kDa
NCBI Accession:	NP_002255
Pathways:	M Phase, Protein targeting to Nucleus

Application Details

Application Notes:	Recombinant human proteins can be used for:
	Native antigens for optimized antibody production
	Positive controls in ELISA and other antibody assays
Comment:	The tag is located at the C-terminal.
Restrictions:	For Research Use only

Handling

50 μg/mL
25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
-80 °C
Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.



Western Blotting

Image 1. Validation with Western Blot