## antibodies -online.com





## **DCXR Protein (Transcript Variant 2) (His tag)**



Image




Go to Product page

| Overview                      |  |
|-------------------------------|--|
| Quantity:                     | 10 μg  |
| Target:                       | DCXR   |
| Protein Characteristics:      | Transcript Variant 2   |
| Origin:                       | Human  |
| Source:                       | Escherichia coli (E. coli)   |
| Protein Type:                 | Recombinant  |
| Purification tag / Conjugate: | This DCXR protein is labelled with His tag.  |
| Application:                  | Antibody Production (AbP), Standard (STD)  |
| Product Details               |  |
| Characteristics:              | <ul> <li>Recombinant human L-xylulose reductase (transcript variant 2) protein expressed in E. coli.</li> <li>Produced with end-sequenced ORF clone</li> </ul>                         |
| Purity:                       | > 95 % as determined by SDS-PAGE and Coomassie blue staining   |
| Endotoxin Level:              | < 0.1 EU per µg protein as determined by LAL test  |
| Target Details                |  |
| Target:                       | DCXR   |
| Alternative Name:             | L-Xylulose Reductase (DCXR Products)   |
| Background:                   | The protein encoded by this gene acts as a homotetramer to catalyze diacetyl reductase and L-xylulose reductase reactions. The encoded protein may play a role in the uronate cycle of |

## **Target Details**

|                     | glucose metabolism and in the cellular osmoregulation in the proximal renal tubules. Defects in this gene are a cause of pentosuria. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug 2010]. |
|---------------------|---|
| Molecular Weight:   | 28.1 kDa  |
| NCBI Accession:     | NP_001182147  |
| Pathways:           | Glycosaminoglycan Metabolic Process, Monocarboxylic Acid Catabolic Process  |
| Application Details |   |
| Application Notes:  | Recombinant human proteins can be used for:   |
|                     | Native antigens for optimized antibody production   |
|                     | Positive controls in ELISA and other antibody assays  |
| Comment:            | The tag is located at the C-terminal.   |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Concentration:      | 50 μg/mL  |
| Buffer:             | 50 mM Tris, 150 mM NaCl, 1 mM DTT, 30 % Glycerol, 1 mM DTT, pH 8.0. Avoid repeated freeze   |
|                     | thaw cycles. Stable for at least 3 months from receipt of products under proper storage and   |
|                     | handling conditions.  |
| Storage:            | -80 °C  |
| Storage Comment:    | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.   |



## **Western Blotting**

Image 1. Validation with Western Blot