# antibodies -online.com





## MAPK14 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image



Go to Product page

$\sim$					
	1//6	r	<b>V</b> I	$\Theta$	Λ

Overview		
Quantity:	20 μg	
Target:	MAPK14	
Protein Characteristics:	Transcript Variant 1	
Origin:	Human	
Source:	HEK-293 Cells	
Protein Type:	Recombinant	
Purification tag / Conjugate:	This MAPK14 protein is labelled with Myc-DYKDDDDK Tag.	
Application:	Antibody Production (AbP), Standard (STD)	
Product Details		
Characteristics:	Recombinant human MAP kinase p38 alpha / MAPK14 (transcript variant 1) protein	
	expressed in HEK293 cells.  • Produced with end-sequenced ORF clone	
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining	
Target Details		
Target:	MAPK14	
Alternative Name:	Map Kinase p38 Alpha,mapk14 (MAPK14 Products)	
Background:	The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as	
	an integration point for multiple biochemical signals, and are involved in a wide variety of	
	cellular processes such as proliferation, differentiation, transcription regulation and	

$\   \text{development. This kinase is activated by various environmental stresses and proinflammatory}$				
cytokines. The activation requires its phosphorylation by MAP kinase kinases (MKKs), or its				
$autophosphorylation\ triggered\ by\ the\ interaction\ of\ MAP3K7IP1/TAB1\ protein\ with\ this\ kinase.$				
The substrates of this kinase include transcription regulator ATF2, MEF2C, and MAX, cell cycle				
regulator CDC25B, and tumor suppressor p53, which suggest the roles of this kinase in stress				
related transcription and cell cycle regulation, as well as in genotoxic stress response. Four				
alternatively spliced transcript variants of this gene encoding distinct isoforms have been				
reported.				

Molecular Weight: 41.3 kDa

NCBI Accession: NP\_001306

Pathways:


MAPK Signaling, Neurotrophin Signaling Pathway, Activation of Innate immune Response,
Cellular Response to Molecule of Bacterial Origin, Regulation of Muscle Cell Differentiation,
Regulation of Cell Size, Hepatitis C, Toll-Like Receptors Cascades, Autophagy, Thromboxane A2
Receptor Signaling, BCR Signaling, S100 Proteins

#### **Application Details**

Application Notes:	Recombinant human proteins can be used for:	
	Native antigens for optimized antibody production	
	Positive controls in ELISA and other antibody assays	
Comment:	The tag is located at the C-terminal.	
Restrictions:	For Research Use only	

#### Handling

Concentration:	50 μg/mL
Buffer:	25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol.
Storage:	-80 °C
Storage Comment: Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze immediately. Only 2-3 freeze thaw cycles are recommended.	



### **Western Blotting**

Image 1. Validation with Western Blot