antibodies - online.com







JNK Protein (Myc-DYKDDDDK Tag)



Image



| _ | | | | | |
|---|---|----|----|---|---|
| U | V | er | VI | е | W |

| Overview | |
|-------------------------------|---|
| Quantity: | 20 μg |
| Target: | JNK (MAPK8) |
| Origin: | Human |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This JNK protein is labelled with Myc-DYKDDDDK Tag. |
| Application: | Antibody Production (AbP), Standard (STD) |
| Product Details | |
| Characteristics: | Recombinant human MAPK8 / JNK1 (transcript variant JNK1-a1) protein expressed in HEK293 cells. |
| | Produced with end-sequenced ORF clone |
| Purity: | > 80 % as determined by SDS-PAGE and Coomassie blue staining |
| Target Details | |
| Target: | JNK (MAPK8) |
| Alternative Name: | Mapk8,jnk1 (MAPK8 Products) |
| Background: | The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as |
| | an integration point for multiple biochemical signals, and are involved in a wide variety of |
| | cellular processes such as proliferation, differentiation, transcription regulation and |
| | development. This kinase is activated by various cell stimuli, and targets specific transcription |

| factors, and thus mediates immediate-early gene expression in response to cell stimuli. The |
|--|
| activation of this kinase by tumor-necrosis factor alpha (TNF-alpha) is found to be required for |
| TNF-alpha induced apoptosis. This kinase is also involved in UV radiation induced apoptosis, |
| which is thought to be related to cytochrom c-mediated cell death pathway. Studies of the |
| mouse counterpart of this gene suggested that this kinase play a key role in T cell proliferation, |
| apoptosis and differentiation. Several alternatively spliced transcript variants encoding distinct |
| isoforms have been reported. |

Receptors Cascades, Signaling of Hepatocyte Growth Factor Receptor, S100 Proteins


| Molecular Weight: | 44 kDa |
|-------------------|--|
| NCBI Accession: | NP_002741 |
| Pathways: | MAPK Signaling, WNT Signaling, TLR Signaling, Fc-epsilon Receptor Signaling Pathway, Neurotrophin Signaling Pathway, Activation of Innate immune Response, Hepatitis C, Toll-Like |

Application Details

| Application Notes: | Recombinant human proteins can be used for: |
|--------------------|--|
| | Native antigens for optimized antibody production |
| | Positive controls in ELISA and other antibody assays |
| Comment: | The tag is located at the C-terminal. |
| Restrictions: | For Research Use only |

Handling

| Concentration: | 50 μg/mL | |
|------------------|---|--|
| Buffer: | 25 mM Tris.HCl, pH 7.3, 100 mM glycine, 10 % glycerol. | |
| Storage: | -80 °C | |
| Storage Comment: | Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze | |
| | immediately. Only 2-3 freeze thaw cycles are recommended. | |



Western Blotting

Image 1. Validation with Western Blot