antibodies .- online.com





MASP1 Protein (Transcript Variant 1) (Myc-DYKDDDDK Tag)



Image

Background:



Go to Product page

Overview	
Quantity:	20 μg
Target:	MASP1
Protein Characteristics:	Transcript Variant 1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This MASP1 protein is labelled with Myc-DYKDDDDK Tag.
Application:	Antibody Production (AbP), Standard (STD)
Product Details	
Characteristics:	 Recombinant human MASP-1 (transcript variant 1) protein expressed in HEK293 cells. Produced with end-sequenced ORF clone
Purity:	> 80 % as determined by SDS-PAGE and Coomassie blue staining
Target Details	
Target:	MASP1
Alternative Name:	Masp-1 (MASP1 Products)

This gene encodes a serine protease that functions as a component of the lectin pathway of

complement activation. The complement pathway plays an essential role in the innate and

adaptive immune response. The encoded protein is synthesized as a zymogen and is activated


when it complexes with the pathogen recognition molecules of lectin pathway, the mannose-

Storage Comment:

anding lectin and the ficolins. This protein is not directly involved in complement activation but ay play a role as an amplifier of complement activation by cleaving complement C2 or by stivating another complement serine protease, MASP-2. The encoded protein is also able to eave fibrinogen and factor XIII and may may be involved in coagulation. A splice variant of this ene which lacks the serine protease domain functions as an inhibitor of the complement athway. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Apr 2010]
) kDa
P_001870
omplement System
ecombinant human proteins can be used for:
ecombinant human proteins can be used for: ative antigens for optimized antibody production
·
ative antigens for optimized antibody production
pative antigens for optimized antibody production positive controls in ELISA and other antibody assays
ative antigens for optimized antibody production ositive controls in ELISA and other antibody assays he tag is located at the C-terminal.
ative antigens for optimized antibody production ositive controls in ELISA and other antibody assays he tag is located at the C-terminal.
eative antigens for optimized antibody production positive controls in ELISA and other antibody assays the tag is located at the C-terminal. For Research Use only
a a transfer and a tr

immediately. Only 2-3 freeze thaw cycles are recommended.

Store at -80°C. Thaw on ice, aliquot to individual single-use tubes, and then re-freeze



Western Blotting

Image 1. Validation with Western Blot